



ANNALES

DE

L'INSTITUT FOURIER

Jean-Louis CLERC & Bent ØRSTED

Conformally invariant trilinear forms on the sphere

Tome 61, n° 5 (2011), p. 1807-1838.

http://aif.cedram.org/item?id=AIF_2011__61_5_1807_0

© Association des Annales de l'institut Fourier, 2011, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

CONFORMALLY INVARIANT TRILINEAR FORMS ON THE SPHERE

by Jean-Louis CLERC & Bent ØRSTED

ABSTRACT. — To each complex number λ is associated a representation π_λ of the conformal group $SO_0(1, n)$ on $\mathcal{C}^\infty(S^{n-1})$ (spherical principal series). For three values $\lambda_1, \lambda_2, \lambda_3$, we construct a trilinear form on $\mathcal{C}^\infty(S^{n-1}) \times \mathcal{C}^\infty(S^{n-1}) \times \mathcal{C}^\infty(S^{n-1})$, which is invariant by $\pi_{\lambda_1} \otimes \pi_{\lambda_2} \otimes \pi_{\lambda_3}$. The trilinear form, first defined for $(\lambda_1, \lambda_2, \lambda_3)$ in an open set of \mathbb{C}^3 is extended meromorphically, with simple poles located in an explicit family of hyperplanes. For generic values of the parameters, we prove uniqueness of trilinear invariant forms.

RÉSUMÉ. — À chaque nombre complexe λ est associée une représentation π_λ du groupe conforme $SO_0(1, n)$ sur $\mathcal{C}^\infty(S^{n-1})$ (série principale sphérique). Pour chaque triplet $(\lambda_1, \lambda_2, \lambda_3)$, nous construisons une forme trilinéaire sur $\mathcal{C}^\infty(S^{n-1}) \times \mathcal{C}^\infty(S^{n-1}) \times \mathcal{C}^\infty(S^{n-1})$ qui est invariante par $\pi_{\lambda_1} \otimes \pi_{\lambda_2} \otimes \pi_{\lambda_3}$. La forme trilinéaire, d'abord définie dans un ouvert de \mathbb{C}^3 est étendue méromorphiquement, avec des pôles simples en une famille explicite de plans de \mathbb{C}^3 . Pour les valeurs génériques des paramètres, nous démontrons l'unicité d'une telle forme trilinéaire invariante.

Introduction

The motivation for this article came from the paper [1] by J. Bernstein and A. Reznikov. In order to estimate automorphic coefficients, they use trilinear invariant forms for $G = PGL_2(\mathbb{R})$. Their paper shows the importance of three related questions of harmonic analysis on $PGL_2(\mathbb{R})$: Given three representations $(\pi_1, \mathcal{H}_1), (\pi_2, \mathcal{H}_2), (\pi_3, \mathcal{H}_3)$ in the principal spherical⁽¹⁾ series of the group G ,

- (i) construct a trilinear invariant functional on $\mathcal{H}_1 \times \mathcal{H}_2 \times \mathcal{H}_3$
- (ii) prove uniqueness (up to a scalar) of such a trilinear invariant functional

Keywords: Trilinear invariant forms, conformal group, meromorphic continuation.
Math. classification: 22E45, 43A85.

⁽¹⁾ with respect to the maximal compact subgroup $K = PO(2)$ of G .

- (iii) compute the value of the trilinear functional on the K fixed vectors of $\mathcal{H}_1, \mathcal{H}_2, \mathcal{H}_3$ respectively.

The representations are realized on the unit circle, on which the group G acts projectively, and indexed by a complex number. One possible generalization consists in replacing the unit circle by the $n - 1$ dimensional sphere S , under the action of the conformal group $G = SO_0(1, n)$. For this case, we present here a construction of an invariant trilinear form (item i)), which uses an analytic continuation over three complex parameters and discuss the uniqueness statement (item ii)) for generic values of the parameters. The computation of the normalization factor (item iii)), even for more geometric situations, will be published elsewhere, (*cf.* [3]).

In Section 1, we recall elementary facts about conformal geometry of the sphere (in particular we give a description of the orbits of G in $S \times S \times S$), and introduce the representations which the paper is concerned with. In Section 2, we define formally the trilinear invariant form, study the domain of convergence of the corresponding integral and determine the analytic continuation in the three complex parameters corresponding to the three representations. In Section 3, we prove the uniqueness statement. The proof relies on Bruhat's theory, which we recall in an appendix at the end of the paper. In Section 4, we give an alternative approach to the construction of an invariant trilinear form, using a realization of the tensor product of two of the representations involved, thus making connection with [12].

The present paper only deals with regular values of the parameters. The residues at poles will yield new conformally invariant trilinear forms, supported by the singular orbits of G in $S \times S \times S$ and involving differential operators akin the Yamabe operator on the sphere, worth of a further study. Other geometric situations are potential domains for similar results. Let P_1, P_2, P_3 be three parabolic subgroups of a semi-simple Lie group, such that G acts on $G/P_1 \times G/P_2 \times G/P_3$ with a finite number of orbits (see [9], [11]). Take three representations induced by characters of P_1, P_2, P_3 . Invariant trilinear forms for these three representations can plausibly be studied along the same lines as in the present paper. The case of three copies of the Shilov boundary S of a bounded symmetric domain of tube type is specially appealing (see [4] for a description of the orbits of G in $S \times S \times S$).

Let us mention the paper by A. Deitmar [5], which has some overlap with our results. Previous work on the subject also includes [13] and [10].

The first author thanks D. Barlet and L. Bérard Bergery for conversations on various aspects of this paper and the Mathematics Department of

Aarhus University for welcoming him during the preparation of the present work.

1. Conformal geometry of the sphere

Let

$$S = \mathbb{S}^{n-1} = \left\{ x = (x_1, x_2, \dots, x_n), \quad |x|^2 = x_1^2 + x_2^2 + \dots + x_n^2 = 1 \right\}$$

be the unit sphere in \mathbb{R}^n . We usually (and tacitly) assume $n \geq 3$, as the case $n = 2$ needs a few minor changes, which are occasionally mentioned in the text. The group $K = SO(n)$ operates on S . Let

$$\mathbf{1}^+ = (1, 0, \dots, 0), \quad \mathbf{1}^- = (-1, 0, \dots, 0).$$

The stabilizer of $\mathbf{1}^+$ in K is the subgroup

$$M \simeq SO(n-1) = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & u \end{pmatrix}, u \in SO(n-1) \right\}.$$

With this notation, $S \simeq K/M$ is a compact Riemannian symmetric space.

Another realization of the sphere is useful. Let $\mathbb{R}^{1,n}$ be the real vector space \mathbb{R}^{n+1} with the quadratic form

$$(1.1) \quad q(x) = [x, x] = x_0^2 - (x_1^2 + x_2^2 + \dots + x_n^2).$$

To $x = (x_1, x_2, \dots, x_n)$ in S associate $\tilde{x} = (1, x_1, x_2, \dots, x_n)$ in $\mathbb{R}^{1,n}$. The correspondance

$$x \longmapsto \mathbb{R}\tilde{x}$$

associates to a point in S an *isotropic line* in $\mathbb{R}^{1,n}$. The correspondance is easily seen to be bijective. The group $G = SO_0(1, n)$ acts naturally on the set of isotropic lines, and hence on S . Explicitly, for x in S and g in G , $g(x)$ is the unique point in S such that

$$\widetilde{g(x)} = (g\tilde{x})_0^{-1} g\tilde{x}.$$

For x in S and g in G , set

$$(1.2) \quad \kappa(g, x) = (g\tilde{x})_0^{-1}.$$

Let x, y be in S . The following identity holds

$$(1.3) \quad [\tilde{x}, \tilde{y}] = 1 - \langle x, y \rangle = \frac{1}{2}|x - y|^2$$

so that for g in G ,

$$(1.4) \quad |g(x) - g(y)| = \kappa(g, x)^{\frac{1}{2}} |x - y| \kappa(g, y)^{\frac{1}{2}}.$$

The infinitesimal version of (1.4) is

$$(1.5) \quad |Dg(x)\xi| = \kappa(g, x)|\xi|$$

for ξ any tangent vector to S at x , where g is in G and $Dg(x)$ stands for the differential at x of the map $x \mapsto g(x)$. Hence the action of G on S is *conformal*, and $\kappa(g, x)$ is interpreted as the *conformal factor* of g at x .

We look at K as a subgroup of G . It is a maximal compact subgroup of G . The stabilizer in G of the point $\mathbf{1}^+$ is the parabolic subgroup $P = MAN$, where

$$A = \left\{ a_t = \begin{pmatrix} \cosh t & \sinh t & 0 & \cdots & 0 \\ \sinh t & \cosh t & 0 & \cdots & 0 \\ 0 & 0 & 1 & & \\ \vdots & \vdots & & \ddots & \\ 0 & 0 & & & 1 \end{pmatrix}, \quad t \in \mathbb{R} \right\}$$

and

$$N = \left\{ n_\xi = \begin{pmatrix} 1 + \frac{|\xi|^2}{2} & -\frac{|\xi|^2}{2} & \xi^t & & \\ \frac{|\xi|^2}{2} & 1 - \frac{|\xi|^2}{2} & \xi^t & & \\ & & 1 & & \\ \xi & -\xi & & \ddots & \\ & & & & 1 \end{pmatrix}, \quad \xi \in \mathbb{R}^{n-1} \right\}.$$

The element a_t ($t \in \mathbb{R}$) acts on S by

$$a_t \begin{pmatrix} x_1 \\ x_2 \\ \cdots \\ x_n \end{pmatrix} = \begin{pmatrix} \frac{\sinh t + x_1 \cosh t}{\cosh t + x_1 \sinh t} \\ \frac{x_2}{\cosh t + x_1 \sinh t} \\ \vdots \\ \frac{x_n}{\cosh t + x_1 \sinh t} \end{pmatrix}.$$

Let \bar{N} be the subgroup image of N by the standard Cartan involution of G ($g \mapsto (g^t)^{-1}$):

$$\bar{N} = \left\{ \bar{n}_\xi = \begin{pmatrix} 1 + \frac{|\xi|^2}{2} & \frac{|\xi|^2}{2} & \xi^t & & \\ -\frac{|\xi|^2}{2} & 1 - \frac{|\xi|^2}{2} & -\xi^t & & \\ & & 1 & & \\ \xi & \xi & & \ddots & \\ & & & & 1 \end{pmatrix}, \quad \xi \in \mathbb{R}^{n-1} \right\}.$$

The map

$$c: \xi \mapsto \bar{n}_\xi(\mathbf{1}^+) = \begin{pmatrix} \frac{1-|\xi|^2}{1+|\xi|^2} \\ \frac{2}{1+|\xi|^2} \xi \end{pmatrix}.$$

is a diffeomorphism from \mathbb{R}^{n-1} onto $S \setminus \{\mathbf{1}^-\}$. Its inverse is the classical *stereographic projection* from the source $\mathbf{1}^-$ onto the tangent space $T_{\mathbf{1}^+}S$ to S at $\mathbf{1}^+$. When using this chart on S , we refer to the *noncompact picture*.

PROPOSITION 1.1. — *The conformal factor $\kappa(g, x)$ is a smooth function of both g and x , which satisfies the following properties:*

- i) $\forall g_1, g_2 \in G, x \in S,$
- $$(1.6) \quad \kappa(g_1 g_2, x) = \kappa(g_1, g_2(x)) \kappa(g_2, x)$$
- ii) $\forall g \in G, x \in S \quad \kappa(g, g^{-1}(x)) = \kappa(g^{-1}, x)^{-1}$
 - iii) $\forall x \in S, k \in K \quad \kappa(k, x) = 1$
 - iv) $\forall x \in S, t \in \mathbb{R}, \quad \kappa(a_t, x) = (\cosh t + x_1 \sinh t)^{-1}.$

Let g in G . As the dimension of the tangent space $T_x S$ is $n - 1$, the Jacobian of g at x is given by

$$(1.7) \quad j(g, x) = \kappa(g, x)^{n-1}.$$

The map $c: \mathbb{R}^{n-1} \rightarrow S \setminus \{-\mathbf{1}\}$ is also conformal. In fact, one has the following relation, valid for any $\xi, \eta \in \mathbb{R}^{n-1}$:

$$(1.8) \quad |c(\xi) - c(\eta)| = \frac{2|\xi - \eta|}{(1 + |\xi|^2)^{\frac{1}{2}}(1 + |\eta|^2)^{\frac{1}{2}}},$$

and its infinitesimal version

$$(1.9) \quad |dc(\xi) \zeta| = \frac{2}{1 + |\xi|^2} |\zeta|$$

($\zeta \in \mathbb{R}^{n-1}$). The corresponding integration formula reads

$$(1.10) \quad \int_S f(x) d\sigma(x) = \int_{\mathbb{R}^{n-1}} f(c(\xi)) \frac{2^{n-1}}{(1 + |\xi|^2)^{n-1}} d\xi.$$

Later, we will need a description of the orbits of G in $S \times S \times S$ (for the diagonal action of G). Recall first that the group G in its diagonal action on $S \times S$ has two orbits:

$$S^2_\top = \{(x, y) \in S \times S, x \neq y\}, \quad \Delta_S = \{(x, x), x \in S\}.$$

As base-point in S^2_\top , choose $(\mathbf{1}^+, \mathbf{1}^-)$. The stabilizer of $(\mathbf{1}^+, \mathbf{1}^-)$ in G is the subgroup MA .

PROPOSITION 1.2. — *Let $n \geq 3$. There are 5 orbits of G in $S \times S \times S$, namely*

$$\begin{aligned} \mathcal{O}_0 &= \{(x_1, x_2, x_3), x_i \neq x_j \text{ for } i \neq j\} \\ \mathcal{O}_1 &= \{(x_1, x, x), x \neq x_1\} \\ \mathcal{O}_2 &= \{(x, x_2, x), x \neq x_2\} \\ \mathcal{O}_3 &= \{(x, x, x_3), x \neq x_3\} \\ \mathcal{O}_4 &= \{(x, x, x)\}. \end{aligned}$$

Proof. — The five subsets of $S \times S \times S$ are invariant under the diagonal action of G . So it suffices to show that G is transitive on each of these sets.

- G is transitive on S , hence on \mathcal{O}_4 . Choose $(\mathbf{1}^+, \mathbf{1}^+, \mathbf{1}^+)$ as base-point. The stabilizer of the base-point in G is the subgroup $P = MAN$.

- The stabilizer P of $\mathbf{1}^+$ is transitive on $S \setminus \{\mathbf{1}^+\}$ (the action of N is already transitive on $S \setminus \{\mathbf{1}^+\}$), so that G is transitive on \mathcal{O}_j , for $j = 1, 2, 3$. As base-point in \mathcal{O}_3 (similar choices can be made for \mathcal{O}_1 and \mathcal{O}_2) choose $(\mathbf{1}^+, \mathbf{1}^+, \mathbf{1}^-)$. The stabilizer of $(\mathbf{1}^+, \mathbf{1}^+, \mathbf{1}^-)$ is the subgroup MA .

- Let x_1, x_2, x_3 be in \mathcal{O}_0 . We may assume w.l.o.g. that $x_1 = \mathbf{1}^+, x_2 = \mathbf{1}^-$ and $x_3 \neq \mathbf{1}^\pm$. The stabilizer of $(\mathbf{1}^+, \mathbf{1}^-)$ in G is MA . The orthogonal projection of x_3 on the hyperplane orthogonal to $(\mathbf{1}^+, \mathbf{1}^-)$ is not 0 and can be mapped by M to ce_2 , with $0 < c < 1$, so that there exists $t \in \mathbb{R}$ such that x_3 is conjugate under M to the point $(\tanh t, \frac{1}{\cosh t}, 0, \dots, 0) = a_t(e_2)$. Hence any triplet in \mathcal{O}_0 is conjugate under G to the triplet $(\mathbf{1}^+, \mathbf{1}^-, e_2)$. Its stabilizer in G is the compact subgroup $\{k \in M, ke_2 = e_2\} \simeq SO(n-2)$. \square

When $n = 2$ (S is the unit circle), then there are two open orbits in $S \times S \times S$ under the action of $SO_0(1, 2)$, each characterized by the value of the *orientation index* of the three points in S . It is possible to remedy to this fact by using the slightly larger (no longer connected) group $O(1, 2)$ instead of $SO_0(1, 2)$.

Let $d\sigma$ be the Lebesgue measure on S and let $\omega_{n-1} = \int_S d\sigma(x)$. Also set $\rho = \frac{n-1}{2}$. Under the action of G , the measure $d\sigma$ is transformed according to

$$(1.11) \quad \int_S f(g^{-1}(x))d\sigma(x) = \int_S f(y)\kappa(g, y)^{2\rho}d\sigma(y)$$

Let λ be in \mathbb{C} . For f in $C^\infty(S)$, the formula

$$(1.12) \quad \pi_\lambda(g)f(x) = \kappa(g^{-1}, x)^{\rho+\lambda} f(g^{-1}(x))$$

defines a *representation* of the group G , which will be denoted by π_λ . It is a continuous representation when the space $\mathcal{C}^\infty(S)$ is equipped with its natural Fréchet topology (see [15] for a systematic study of these representations).

The representations π_λ and $\pi_{-\lambda}$ are dual in the sense that, for all $\varphi, \psi \in \mathcal{C}^\infty(S)$

$$(1.13) \quad \int_S \pi_{-\lambda}(g)\varphi(s)\pi_\lambda(g)\psi(s) ds = \int_S \varphi(s)\psi(s) ds$$

as can be deduced from the change of variable formula (1.11). For λ *pure imaginary*, the representation π_λ can be extended continuously to $L^2(S)$ to yield a *unitary* representation of G (this is the reason for using $\rho + \lambda$ in the definition of π_λ). Observe that the action of K is independent of λ and the constant function $\mathbb{1}_S$ (equal to 1 everywhere) is fixed by the action of K .

For α in \mathbb{C} , let k_α be the kernel on $S \times S$ defined by

$$k_\alpha(x, y) = |x - y|^{-\rho+\alpha}.$$

It satisfies the following *transformation property* under the action of an element g of G :

$$(1.14) \quad k_\alpha(g(x), g(y)) = \kappa(g, x)^{-\frac{\rho}{2}+\frac{\alpha}{2}} k_\alpha(x, y) \kappa(g, y)^{-\frac{\rho}{2}+\frac{\alpha}{2}}$$

for all x, y in S .

2. Construction of an invariant trilinear form

2.1. Formal construction

Let $\lambda_1, \lambda_2, \lambda_3$ be three complex numbers. Let \mathcal{T} be a continuous trilinear form from $\mathcal{C}^\infty(S) \times \mathcal{C}^\infty(S) \times \mathcal{C}^\infty(S)$ into \mathbb{C} . The functional \mathcal{T} is said to be *invariant* w.r.t. $\pi_{\lambda_1}, \pi_{\lambda_2}, \pi_{\lambda_3}$ if, for every f_1, f_2, f_3 in $\mathcal{C}^\infty(S)$, and every g in G ,

$$(2.1) \quad \mathcal{T}(\pi_{\lambda_1}(g)f_1, \pi_{\lambda_2}(g)f_2, \pi_{\lambda_3}(g)f_3) = \mathcal{T}(f_1, f_2, f_3).$$

By Schwartz's *kernel theorem*, there exists a unique distribution T on $S \times S \times S$, such that

$$(2.2) \quad \mathcal{T}(f_1, f_2, f_3) = T(f_1 \otimes f_2 \otimes f_3)$$

where, as usual, $f_1 \otimes f_2 \otimes f_3$ is the function on $S \times S \times S$ defined by

$$(f_1 \otimes f_2 \otimes f_3)(x_1, x_2, x_3) = f_1(x_1)f_2(x_2)f_3(x_3).$$

Let $\alpha_1, \alpha_2, \alpha_3$ be complex numbers, and set $\alpha = (\alpha_1, \alpha_2, \alpha_3)$. Let K_α be the kernel on $S \times S \times S$ defined by

$$(2.3) \quad K_\alpha(x_1, x_2, x_3) = k_{\alpha_1}(x_2, x_3) k_{\alpha_2}(x_3, x_1) k_{\alpha_3}(x_1, x_2)$$

For f_1, f_2, f_3 three functions in $C^\infty(S)$, define the trilinear functional \mathcal{K}_α by

$$(2.4) \quad \mathcal{K}_\alpha(f_1, f_2, f_3) = \int_{S \times S \times S} K_\alpha(x_1, x_2, x_3) f_1(x_1) f_2(x_2) f_3(x_3) \times d\sigma(x_1) d\sigma(x_2) d\sigma(x_3)$$

whenever it makes sense.

THEOREM 2.1. — *Let $\lambda = (\lambda_1, \lambda_2, \lambda_3) \in \mathbb{C}^3$. Define $\alpha = (\alpha_1, \alpha_2, \alpha_3)$ by the relations*

$$(2.5) \quad \begin{aligned} \alpha_1 &= -\lambda_1 + \lambda_2 + \lambda_3 \\ \alpha_2 &= -\lambda_2 + \lambda_3 + \lambda_1 \\ \alpha_3 &= -\lambda_3 + \lambda_1 + \lambda_2. \end{aligned}$$

Then

$$(2.6) \quad \mathcal{K}_\alpha(\pi_{\lambda_1}(g)f_1, \pi_{\lambda_2}(g)f_2, \pi_{\lambda_3}(g)f_3) = \mathcal{K}_\alpha(f_1, f_2, f_3),$$

whenever the integral on the right handside is defined.

The proof is obtained by the change of variables $y_j = g^{-1}(x_j)$ ($j = 1, 2, 3$) in the integral defining the left hand-side, using (1.14) and (1.11). Observe that the right-handside integral converges if, for $i \neq j$, the supports of f_i and f_j are disjoint, or if

$$\int_{S \times S \times S} |K_\alpha(x_1, x_2, x_3)| d\sigma(x_1) d\sigma(x_2) d\sigma(x_3) < +\infty.$$

The version of the trilinear functional in the noncompact picture will be useful. Let $\alpha = (\alpha_1, \alpha_2, \alpha_3) \in \mathbb{C}^3$ and set for $y_1, y_2, y_3 \in \mathbb{R}^{n-1}$

$$(2.7) \quad J_\alpha(y_1, y_2, y_3) = |y_1 - y_2|^{-\rho+\alpha_3} |y_2 - y_3|^{-\rho+\alpha_1} |y_3 - y_1|^{-\rho+\alpha_2}$$

and, whenever it makes sense, let \mathcal{J}_α be the associated distribution on $\mathbb{R}^{n-1} \times \mathbb{R}^{n-1} \times \mathbb{R}^{n-1}$ given by

$$(2.8) \quad \mathcal{J}_\alpha(\varphi) = \int J_\alpha(y_1, y_2, y_3) \varphi(y_1, y_2, y_3) dy_1 dy_2 dy_3,$$

($\varphi \in \mathcal{C}^\infty(\mathbb{R}^{n-1} \times \mathbb{R}^{n-1} \times \mathbb{R}^{n-1})$). Moreover, let Ψ_α be the function defined on $\mathbb{R}^{n-1} \times \mathbb{R}^{n-1} \times \mathbb{R}^{n-1}$ by

$$\Psi_\alpha(y_1, y_2, y_3) = (1 + |y_1|^2)^{-\rho - \frac{\alpha_2}{2} - \frac{\alpha_3}{2}} (1 + |y_2|^2)^{-\rho - \frac{\alpha_3}{2} - \frac{\alpha_1}{2}} \times (1 + |y_3|^2)^{-\rho - \frac{\alpha_1}{2} - \frac{\alpha_2}{2}}.$$

PROPOSITION 2.2. — *Let $f \in \mathcal{C}^\infty(S \times S \times S)$. Then*

$$(2.9) \quad \mathcal{K}_\alpha(f) = 2^{3(n-1)} \mathcal{J}_\alpha((f \circ c)\Psi_\alpha)$$

whenever the left hand side is defined.

This is merely the change of variable (1.10) in the integral (2.4).

2.2. Integrability of the kernel K_α

THEOREM 2.3. — *The kernel K_α is integrable on $S \times S \times S$ if and only if*

$$(2.10) \quad \Re\alpha_j > -\rho, \quad j = 1, 2, 3$$

$$(2.11) \quad \Re\alpha_1 + \Re\alpha_2 + \Re\alpha_3 > -\rho.$$

Proof. — It is enough to study the integral when the α 's are real, in which case the kernel K_α is positive. Let \mathcal{U} be a (small) neighborhood of $(\mathbf{1}^+, \mathbf{1}^+, \mathbf{1}^+)$ in $S \times S \times S$. Let g be in G . As x varies in S , $j(g, x)$ remains bounded from below and from above. Thanks to the transformation property of the kernel k_α (1.14), the integrals of K_α over \mathcal{U} and over $g(\mathcal{U})$ are of the same nature. As \mathcal{U} meets all G -orbits, a partition of unity argument shows that the integrability over S of the kernel K_α is equivalent to its integrability over \mathcal{U} . Now use the stereographic projection to see that the integrability of K_α over \mathcal{U} is equivalent to the integrability of J_α over $c^{-1}(\mathcal{U})$, which is a (small) neighborhood of $(0, 0, 0)$ in $\mathbb{R}^{n-1} \times \mathbb{R}^{n-1} \times \mathbb{R}^{n-1}$.

Hence, we are reduced to study the convergence of the integral

$$I_1 = \int_{\substack{|\xi_1| < \delta \\ |\xi_2| < \delta \\ |\xi_3| < \delta}} |\xi_1 - \xi_2|^{\alpha_3 - \rho} |\xi_2 - \xi_3|^{\alpha_1 - \rho} |\xi_3 - \xi_1|^{\alpha_2 - \rho} d\xi_1 d\xi_2 d\xi_3,$$

where δ is a small positive number. Set

$$y_1 = \xi_1, \quad y_2 = \xi_1 - \xi_2, \quad y_3 = -\xi_2 + \xi_3.$$

Then, after integrating with respect to y_1 , the integral I_1 is seen to be of the same nature as the integral

$$I_2 = \int_{\substack{|y_2| < \delta \\ |y_3| < \delta}} |y_2|^{\alpha_3 - \rho} |y_3|^{\alpha_2 - \rho} |y_2 - y_3|^{\alpha_1 - \rho} dy_2 dy_3.$$

Let Σ be the unit ball in $\mathbb{R}^{(n-1)} \times \mathbb{R}^{(n-1)}$, and let $d\sigma$ be the Lebesgue measure on Σ . Use polar coordinates $(r, (\sigma_2, \sigma_3))$ defined by

$$y_2 = r\sigma_2, \quad y_3 = r\sigma_3, \quad r^2 = |y_2|^2 + |y_3|^2, \quad |\sigma_1|^2 + |\sigma_3|^2 = 1$$

to obtain that I_2 is of the same nature as

$$I_3 = \int_0^\delta r^{\alpha_1 + \alpha_2 + \alpha_3 - 3\rho + 2(n-1) - 1} dr \times \int_\Sigma |\sigma_2|^{\alpha_3 - \rho} |\sigma_3|^{\alpha_2 - \rho} |\sigma_2 - \sigma_3|^{\alpha_1 - \rho} d\sigma.$$

The first factor converges if and only if condition (2.11) is satisfied. It remains to discuss the convergence of

$$I_4 = \int_\Sigma |\sigma_2|^{\alpha_3 - \rho} |\sigma_3|^{\alpha_2 - \rho} |\sigma_2 - \sigma_3|^{\alpha_1 - \rho} d\sigma.$$

Let $\delta > 0$ and consider the following open subsets of Σ :

$$\begin{aligned} \Sigma_2 &= \{(\sigma_2, \sigma_3), |\sigma_2| < \delta\}, \\ \Sigma_3 &= \{(\sigma_2, \sigma_3), |\sigma_3| < \delta\}, \\ \Sigma_1 &= \{(\sigma_2, \sigma_3), |\sigma_2 - \sigma_3| < \delta\}. \end{aligned}$$

For δ small enough, these sets are disjoint (recall that $|\sigma_2|^2 + |\sigma_3|^2 = 1$). On $\Sigma \setminus (\Sigma_1 \cup \Sigma_2 \cup \Sigma_3)$ the function to be integrated is bounded from below. Hence the integral I_4 is convergent if and only the integrals

$$K_j = \int_{\Sigma_j} |\sigma_2|^{\alpha_3 - \rho} |\sigma_3|^{\alpha_2 - \rho} |\sigma_2 - \sigma_3|^{\alpha_1 - \rho} d\sigma$$

are convergent for $j = 1, 2, 3$. Let $j = 2$. On Σ_2 , both $|\sigma_3|$ and $|\sigma_2 - \sigma_3|$ are bounded from below, so that it is equivalent to study the convergence of the integral

$$\int_{\Sigma_2} |\sigma_2|^{\alpha_3 - \rho} d\sigma.$$

We are reduced to a classical situation and K_2 converges if and only if $\alpha_3 - \rho > -(n - 1)$. A similar study applies to K_1 and K_3 . This completes the proof of Theorem 2.3. □

COROLLARY 2.4. — *Let $\lambda_1, \lambda_2, \lambda_3$ be three complex numbers satisfying the condition*

$$0 \leq \Re(\lambda_j) < \rho.$$

Define α by the relations (2.5). Then the kernel K_α is integrable, and the corresponding trilinear form is invariant for $\pi_{\lambda_1}, \pi_{\lambda_2}, \pi_{\lambda_3}$.

Proof. — The conditions on $\lambda = (\lambda_1, \lambda_2, \lambda_3)$ imply that $\Re(\alpha_j) > -\rho$ for $j = 1, 2, 3$ and $\Re(\alpha_1 + \alpha_2 + \alpha_3) \geq 0$. Hence α is in the domain of integrability of K_α . □

The corollary covers all interesting cases for spherical unitary series, provided one excludes the trivial representation. In fact, the parameter λ for such a representation can be chosen either as pure imaginary (principal series) or satisfying $0 < \lambda < \rho$ (complementary series, excluding the trivial representation).

2.3. Analytic continuation of \mathcal{K}_α

The main result of this section concerns the analytic continuation of \mathcal{K}_α beyond its domain of integrability.

THEOREM 2.5. — *The map $\alpha \mapsto \mathcal{K}_\alpha$, originally defined for α in the domain described by the conditions (2.10) and (2.11), valued in $\mathcal{D}'(S \times S \times S)$ can be extended meromorphically with at most simple poles along the family of hyperplanes in $\mathbb{C} \times \mathbb{C} \times \mathbb{C}$ defined by the equations*

$$(2.12) \quad \alpha_j = -\rho - 2k, \quad j = 1, 2, 3, \quad k \in \mathbb{N}$$

$$(2.13) \quad \alpha_1 + \alpha_2 + \alpha_3 = -\rho - 2k, \quad k \in \mathbb{N}.$$

Proof. — Let f be in $\mathcal{C}^\infty(S \times S \times S)$ and consider the integral

$$\int_{S \times S \times S} K_\alpha(x_1, x_2, x_3) f(x_1, x_2, x_3) d\sigma(x_1) d\sigma(x_2) d\sigma(x_3).$$

to be meromorphically continued. Repeating the argument given supra during the discussion of the integrability of the kernel K_α , we may assume that f has its support contained in a small neighborhood of the point $(\mathbf{1}^+, \mathbf{1}^+, \mathbf{1}^+)$. Further, transfer the integral to the noncompact picture (cf. (2.9)), and study the analytic continuation of

$$\alpha \mapsto \mathcal{J}_\alpha((f \circ c)\Psi_\alpha)$$

Now $\varphi = f \circ c$ is in $\mathcal{C}_c^\infty(\mathbb{R}^{n-1} \times \mathbb{R}^{n-1} \times \mathbb{R}^{n-1})$, and both $\alpha \mapsto \Psi_\alpha$ and $\alpha \mapsto \Psi_\alpha^{-1}$ are entire functions on \mathbb{C}^3 , so that it is equivalent to study the meromorphic continuation of \mathcal{J}_α as a distribution on $\mathbb{R}^{n-1} \times \mathbb{R}^{n-1} \times \mathbb{R}^{n-1}$.

The kernel J_α is invariant by translations by “diagonal vectors”. To take advantage of this remark, make the change of variables

$$z_1 = y_1, \quad z_2 = y_1 - y_3, \quad z_3 = y_1 - y_2$$

in the integral

$$\begin{aligned} \mathcal{J}_\alpha(\varphi) = \int & |y_1 - y_2|^{-\rho+\alpha_3} |y_2 - y_3|^{-\rho+\alpha_1} |y_3 - y_1|^{-\rho+\alpha_2} \\ & \times \varphi(y_1, y_2, y_3) dy_1 dy_2 dy_3 \end{aligned}$$

to obtain

$$\mathcal{I}_\alpha(\varphi) = \int_{\mathbb{R}^{n-1} \times \mathbb{R}^{n-1}} |z_2|^{-\rho+\alpha_2} |z_3|^{-\rho+\alpha_3} |z_2 - z_3|^{-\rho+\alpha_1} \psi(z_2, z_3) dz_2 dz_3$$

where we have set

$$\psi(z_2, z_3) = \int_{\mathbb{R}^{n-1}} \varphi(z_1, z_1 - z_3, z_1 - z_2) dz_1.$$

Now observe that ψ is in $C_c^\infty(\mathbb{R}^{n-1} \times \mathbb{R}^{n-1})$. Hence we are reduced to studying the analytic continuation of the distribution \mathcal{I}_α on $\mathbb{R}^{n-1} \times \mathbb{R}^{n-1}$ defined by

$$(2.14) \quad \mathcal{I}_\alpha(\psi) = \int_{\mathbb{R}^{n-1} \times \mathbb{R}^{n-1}} I_\alpha(z_2, z_3) \psi(z_2, z_3) dz_2 dz_3$$

for ψ in $C_c^\infty(\mathbb{R}^{n-1} \times \mathbb{R}^{n-1})$, where we set

$$(2.15) \quad I_\alpha(z_2, z_3) = |z_2|^{-\rho+\alpha_2} |z_3|^{-\rho+\alpha_3} |z_2 - z_3|^{-\rho+\alpha_1}.$$

For $\delta > 0$, consider the following open subsets of $\mathbb{R}^{n-1} \times \mathbb{R}^{n-1}$

$$\begin{aligned} \mathcal{U}_0 &= \left\{ (z_2, z_3), |z_2| < \delta, |z_3| < \delta, |z_2 - z_3| < \delta \right\} \\ \mathcal{U}_1 &= \left\{ (z_2, z_3), |z_2| > \frac{\delta}{2}, |z_3| > \frac{\delta}{2}, |z_2 - z_3| < \frac{\delta}{2} \right\} \\ \mathcal{U}_2 &= \left\{ (z_2, z_3), |z_2| < \frac{\delta}{2}, |z_3| > \frac{\delta}{2}, |z_3 - z_2| > \frac{\delta}{2} \right\} \\ \mathcal{U}_3 &= \left\{ (z_2, z_3), |z_2| > \frac{\delta}{2}, |z_3| < \frac{\delta}{2}, |z_2 - z_3| > \frac{\delta}{2} \right\} \\ \mathcal{U}_\infty &= \left\{ (z_2, z_3), |z_2| > \frac{\delta}{2}, |z_3| > \frac{\delta}{2}, |z_2 - z_3| > \frac{\delta}{2} \right\}. \end{aligned}$$

The family of these five open sets form a covering of $\mathbb{R}^{n-1} \times \mathbb{R}^{n-1}$. Let study the restriction of the distribution \mathcal{I}_α to each of these five open subsets.

If $\text{Supp}(\psi) \subset \mathcal{U}_\infty$, then the $\mathcal{I}_\alpha(\psi)$ extends as an entire function, because I_α has no singularity on \mathcal{U}_∞ . Next assume that $\text{Supp}(\psi) \subset \mathcal{U}_2$. Set

$$\phi_\alpha(z_2) = \int_{\mathbb{R}^{n-1}} |z_3|^{-\rho+\alpha_3} |z_2 - z_3|^{-\rho+\alpha_1} \psi(z_2, z_3) dz_3,$$

so that

$$(2.16) \quad \mathcal{I}_\alpha(\psi) = \int_{\mathbb{R}^{n-1}} |z_2|^{-\rho+\alpha_2} \phi_\alpha(z_2) dz_2.$$

As $|z_3|$ and $|z_2 - z_3|$ are bounded from below on \mathcal{U}_2 , the function ϕ_α is in $C_c^\infty(\mathbb{R}^{n-1})$ and $\alpha \mapsto \phi_\alpha$ is an entire function on \mathbb{C}^3 . On \mathbb{R}^{n-1} , the distribution-valued function $s \mapsto |z_2|^s$ extends meromorphically on \mathbb{C} ,

with simple poles at $s = -(n - 1) - 2k, k \in \mathbb{N}$ (see [6]), so that the integral (2.16) extends meromorphically to \mathbb{C}^3 with simple poles along the hyperplanes $\alpha_2 = -\rho - 2k$. A similar analysis can be done over \mathcal{U}_1 and \mathcal{U}_3 .

To sum up what we have already proved, introduce the family \mathcal{M} of meromorphic functions on \mathbb{C}^3 having at most simple poles along the hyperplanes $\{\alpha_j = -\rho - 2k\}, j = 1, 2, 3, k \in \mathbb{N}$. Notice that they are the hyperplanes corresponding to the conditions (2.12).

PROPOSITION 2.6. — *Let ψ be in $C_c^\infty(\mathbb{R}^{n-1} \times \mathbb{R}^{n-1})$ and assume that $\text{Supp}(\psi) \not\ni (0, 0)$. Then the function $\alpha \mapsto \mathcal{I}_\alpha(\psi)$ belongs to the class \mathcal{M} .*

Proof. — In fact, choose δ small enough so that $\text{Supp}(\psi) \cap \mathcal{U}_0 = \emptyset$. Use a partition of unity to write ψ as

$$\psi = \psi_1 + \psi_2 + \psi_3 + \psi_\infty$$

where $\psi_j \in C_c^\infty(\mathbb{R}^{n-1} \times \mathbb{R}^{n-1})$ and $\text{Supp}(\psi_j) \subset \mathcal{U}_j, j = 1, 2, 3, \infty$. The previous analysis shows that $\alpha \mapsto \mathcal{I}_\alpha(\psi_j)$ extends meromorphically on \mathbb{C}^3 with at most simple poles along the hyperplanes $\alpha_j = -\rho - 2k, k \in \mathbb{N}$ for $j = 1, 2, 3$, whereas $\mathcal{I}_\alpha(\psi_\infty)$ is an entire function of α . \square

Now we use *a priori* the existence of the meromorphic continuation of such integrals (see [14]). Moreover, the poles are located on a locally finite family of affine hyperplanes (of a rather specific type, but we won't need this result). Let \mathcal{H} be such an hyperplane (to be determined), but not included in the family of hyperplanes given by conditions (2.12). Let $\alpha^0 = (\alpha_1^0, \alpha_2^0, \alpha_3^0)$ be a regular point in \mathcal{H} , (*i.e.* not contained in any other hyperplane of poles). The Laurent coefficients at α^0 are distributions on $\mathbb{R}^{n-1} \times \mathbb{R}^{n-1}$, and, by Lemma 2.6, their supports have to be contained in $\{0, 0\}$, hence they are derivatives of the Dirac measure $\delta_{(0,0)}$. So, if \mathcal{I}_α does have a pole at α^0 , there exists a smooth function ρ on $\mathbb{R}^{n-1} \times \mathbb{R}^{n-1}$ with compact support and identically equal to 1 in a neighbourhood of $(0, 0)$, and a polynomial P on $\mathbb{R}^{n-1} \times \mathbb{R}^{n-1}$, homogeneous of degree k such that $\mathcal{I}_\alpha(\rho P)$ does not extend holomorphically at α^0 . For $t \in \mathbb{R}^*$, let $\rho_t(z_2, z_3) = \rho(tz_2, tz_3)$. Use the change of variables $(z_2 \mapsto tz_2, z_3 \mapsto tz_3)$ to get ⁽²⁾, for $\alpha \neq \alpha_0$

$$\begin{aligned} \mathcal{I}_\alpha(\rho_t P) &= \int_{\mathbb{R}^{n-1} \times \mathbb{R}^{n-1}} |z_2|^{-\rho+\alpha_3} |z_3|^{-\rho+\alpha_2} |z_2 - z_3|^{-\rho+\alpha_1} \\ &\quad \times \rho(tz_2, tz_3) P(z_2, z_3) dz_2 dz_3 \\ &= |t|^{-\alpha_3-\alpha_2-\alpha_1-\rho} t^{-k} \mathcal{I}_\alpha(\rho P). \end{aligned}$$

⁽²⁾ Following a traditional way, we write $\mathcal{I}_\alpha(\varphi)$ as an integral. What is really used here is merely the homogeneity of \mathcal{I}_α .

Hence

$$(2.17) \quad (1 - |t|^{-\alpha_3 - \alpha_2 - \alpha_1 - \rho} t^{-k}) \mathcal{I}_\alpha(\rho P) = \mathcal{I}_\alpha((\rho - \rho_t)P).$$

Now observe that the support of $(\rho - \rho_t)P$ does not contain $(0, 0)$, and hence, $\alpha \mapsto \mathcal{I}_\alpha((\rho - \rho_t)P)$ belongs to \mathcal{M} . The assumption that $\mathcal{I}_\alpha(\rho P)$ does have a pole at α_0 forces the condition

$$\forall t \in \mathbb{R}^*, \quad 1 - |t|^{-\alpha_3^0 - \alpha_2^0 - \alpha_1^0 - \rho} t^{-k} = 0.$$

In turn, this condition amounts to

$$k \in 2\mathbb{N}, \quad \alpha_1^0 + \alpha_2^0 + \alpha_3^0 = -\rho - k.$$

Moreover, (2.17) shows that \mathcal{I}_α has at most a simple pole along the corresponding hyperplane. This achieves the proof of Theorem 2.5. \square

As the invariance condition remains true by analytic continuation, Theorem 2.5 can be reformulated for trilinear invariant functionals (cf. Theorem 2.1).

THEOREM 2.7. — *Let $\lambda = (\lambda_1, \lambda_2, \lambda_3)$ in \mathbb{C}^3 . Assume that*

$$(2.18) \quad \begin{aligned} -\lambda_1 + \lambda_2 + \lambda_3 &\notin -\rho - 2\mathbb{N} \\ \lambda_1 - \lambda_2 + \lambda_3 &\notin -\rho - 2\mathbb{N} \\ \lambda_1 + \lambda_2 - \lambda_3 &\notin -\rho - 2\mathbb{N} \\ \lambda_1 + \lambda_2 + \lambda_3 &\notin -\rho - 2\mathbb{N}. \end{aligned}$$

Set $\alpha = (\alpha_1, \alpha_2, \alpha_3)$ where (cf. (2.5))

$$\begin{aligned} \alpha_1 &= -\lambda_1 + \lambda_2 + \lambda_3 \\ \alpha_2 &= -\lambda_2 + \lambda_3 + \lambda_1 \\ \alpha_3 &= -\lambda_3 + \lambda_1 + \lambda_2. \end{aligned}$$

Then $(f_1, f_2, f_3) \mapsto \mathcal{T}_\lambda(f_1, f_2, f_3) = \mathcal{K}_\alpha(f_1 \otimes f_2 \otimes f_3)$ is a well defined non trivial trilinear invariant functional w.r.t. the representations $(\pi_{\lambda_1}, \pi_{\lambda_2}, \pi_{\lambda_3})$.

The next result was obtained some time ago by the present authors, and has been generalized to other geometric situations in a collaboration with T. Kobayashi and M. Pevzner (see [3]). To state the result, consider the evaluation of the functional \mathcal{K}_α against the function $\mathbb{I}_S \otimes \mathbb{I}_S \otimes \mathbb{I}_S$, where \mathbb{I}_S is the function which is identically 1 on S . Let

$$\begin{aligned} I(\lambda) = K(\alpha) &= \int_S \int_S \int_S k_{\alpha_1}(x_2, x_3) k_{\alpha_2}(x_3, x_1) k_{\alpha_3}(x_1, x_2) \\ &\quad \times d\sigma(x_1) d\sigma(x_2) d\sigma(x_3) \end{aligned}$$

where α and λ are related by the relations (2.5).

PROPOSITION 2.8. — *Let $\lambda = (\lambda_1, \lambda_2, \lambda_3)$ in \mathbb{C}^3 and assume that the conditions (2.18) are satisfied. Then*

$$(2.19) \quad I(\lambda) = \left(\frac{\sqrt{\pi}}{2}\right)^{3(n-1)} 2^{\lambda_1+\lambda_2+\lambda_3} \frac{\Gamma\left(\frac{\lambda_1+\lambda_2+\lambda_3+\rho}{2}\right)\Gamma\left(\frac{-\lambda_1+\lambda_2+\lambda_3+\rho}{2}\right)\Gamma\left(\frac{\lambda_1-\lambda_2+\lambda_3+\rho}{2}\right)\Gamma\left(\frac{\lambda_1+\lambda_2-\lambda_3+\rho}{2}\right)}{\Gamma(\rho + \lambda_1)\Gamma(\rho + \lambda_2)\Gamma(\rho + \lambda_3)}.$$

Remark. — Both sides of the formula are meromorphic functions on \mathbb{C}^3 , and they are equal where defined. Notice that $I(\lambda) = K(\alpha)$ has simple poles exactly as prescribed by Theorem 2.5.

This result allows to strengthen the previous theorem. Define

$$\tilde{\mathcal{K}}_\alpha = \frac{\mathcal{K}_\alpha}{\Gamma\left(\frac{\lambda_1+\lambda_2+\lambda_3+\rho}{2}\right)\Gamma\left(\frac{-\lambda_1+\lambda_2+\lambda_3+\rho}{2}\right)\Gamma\left(\frac{\lambda_1-\lambda_2+\lambda_3+\rho}{2}\right)\Gamma\left(\frac{\lambda_1+\lambda_2-\lambda_3+\rho}{2}\right)}.$$

THEOREM 2.9. — *The distribution-valued function $\alpha \mapsto \tilde{\mathcal{K}}_\alpha$ extends as an entire holomorphic function on \mathbb{C}^3 . The trilinear functional $\tilde{\mathcal{T}}_\lambda$ defined by*

$$\tilde{\mathcal{T}}_\lambda(f_1, f_2, f_3) = \tilde{\mathcal{K}}_\alpha(f_1 \otimes f_2 \otimes f_3)$$

on $\mathcal{C}^\infty(S) \times \mathcal{C}^\infty(S) \times \mathcal{C}^\infty(S)$ is invariant with respect to $(\pi_{\lambda_1}, \pi_{\lambda_2}, \pi_{\lambda_3})$, where $(\lambda_1, \lambda_2, \lambda_3)$ are related to $(\alpha_1, \alpha_2, \alpha_3)$ by the relations (2.5). The trilinear form $\tilde{\mathcal{T}}_\lambda$ is not identically 0 provided the two following conditions are not simultaneously realized

- $\exists j, 1 \leq j \leq 3, \quad \lambda_j \in -\rho - \mathbb{N}$
- (at least) one of the conditions (2.18) is satisfied .

Proof. — The function $\alpha \mapsto \tilde{\mathcal{K}}_\alpha$ extends holomorphically near any regular point of the hyperplanes defined by conditions (2.12) and (2.13), so is holomorphic outside of the set where at least two hyperplanes of poles meet. But this set is of codimension 2 and hence $\alpha \mapsto \tilde{\mathcal{K}}_\alpha$ extends as a holomorphic function to all of \mathbb{C}^3 . If none of the conditions (2.18) is satisfied, then $\tilde{\mathcal{K}}_\alpha$ is a multiple ($\neq 0$) of \mathcal{K}_α which is different from 0 on Ω . If $\lambda_j \notin -\rho + \mathbb{N}$ for $j = 1, 2, 3$, then $I(\lambda) \neq 0$ and hence \mathcal{K}_α is not identically 0. □

3. Uniqueness of the invariant trilinear form

3.1. Induced representations and line bundles

For this part, it is useful to realize the representation π_λ as acting on smooth sections of a line bundle over S .

Recall that the stabilizer of $\mathbf{1}^+$ in G is the parabolic subgroup $P = MAN$. The left invariant Haar measure on P is

$$(3.1) \quad \int_P f(p)dp = \int_M \int_A \int_N f(man) dm da dn.$$

For $\lambda \in \mathbb{C}$, denote by χ_λ the character of P defined by

$$\chi_\lambda(man) = e^{t\lambda}.$$

The modular function of P is given by $\delta_P(man) = e^{-2\rho(\log a)} = \chi_{-2\rho}(p)$ (see e.g. [18] Lemma 5.5.1.1), so that, for any q in P

$$(3.2) \quad \int_P f(pq) dp = \chi_{2\rho}(q) \int_P f(p) dp.$$

Let \mathcal{E}_λ be the space of functions f in $C^\infty(G)$ which, for all g in G , p in P satisfy

$$(3.3) \quad f(gp) = \chi_{-(\rho+\lambda)}(p)f(g).$$

Then G acts on \mathcal{E}_λ by

$$(3.4) \quad \Pi_\lambda(g)f(\gamma) = f(g^{-1}\gamma).$$

To any function f in \mathcal{E}_λ , associate the function f^\sim defined on S by the formula

$$(3.5) \quad f^\sim(s) = f(k)$$

where k is any element in K satisfying $k\mathbf{1}^+ = s$. As $f(km) = f(m)$ for any m in M , the right handside of (3.5) does not depend on m , but merely on $k\mathbf{1}^+ = s$.

Now, let g be in G , let s be in S , and let k be in K such that $k\mathbf{1}^+ = s \in S$. Let

$$g^{-1}k = k(g^{-1}k) a(g^{-1}k) n$$

be the Iwasawa decomposition of $g^{-1}k$. Then

$$k(g^{-1}k)\mathbf{1}^+ = (g^{-1}k)(\mathbf{1}^+) = g^{-1}(s)$$

and

$$\kappa(g^{-1}, s) = \kappa(g^{-1}k, \mathbf{1}^+) = \chi_{-1}(a(g^{-1}k)).$$

Hence,

$$\begin{aligned} (\Pi_\lambda(g)f)^\sim(s) &= f(g^{-1}k) = f(k(g^{-1}k) a(g^{-1}k) n) \\ &= \kappa(g^{-1}, s)^{\rho+\lambda} f^\sim(g^{-1}(s)) \end{aligned}$$

so that $f \mapsto f^\sim$ is an intertwining operator for Π_λ and π_λ .

Let χ be a character of A . Denote by \mathbb{C}_χ the representation of P on \mathbb{C} given by

$$ma \cdot z = \chi(a)z.$$

Form the line bundle $L = L_\chi = G \times_P \mathbb{C}_\chi$ over S , and let $\mathcal{L} = \mathcal{L}_\chi^\infty$ be the space of \mathcal{C}^∞ sections of L . Then G acts naturally on \mathcal{L} by left translations. As sections of L can be identified with functions on G transforming by χ^{-1} under the right action of P , the representation of G on \mathcal{L} is equivalent to π_λ if $\chi = \chi_{\rho+\lambda}$.

Now take three characters $\chi_1 = \chi_{\rho+\lambda_1}, \chi_2 = \chi_{\rho+\lambda_2}, \chi_3 = \chi_{\rho+\lambda_3}$ of A , set $\lambda = (\lambda_1, \lambda_2, \lambda_3)$ and form the “exterior” product bundle $L_\lambda = L_{\chi_1} \boxtimes L_{\chi_2} \boxtimes L_{\chi_3}$ as a line bundle over $S \times S \times S$. Let \mathcal{L}_λ be the space of \mathcal{C}^∞ sections of this bundle. Let the group G acts naturally on this space by diagonal action. Then a trilinear invariant functional on $\mathcal{C}^\infty(S) \times \mathcal{C}^\infty(S) \times \mathcal{C}^\infty(S)$ for $\pi_{\lambda_1}, \pi_{\lambda_2}, \pi_{\lambda_3}$ corresponds to an invariant linear form on \mathcal{L}_λ .

In turn, such an invariant linear functional on \mathcal{L}_λ can be viewed as an invariant distribution density for the dual bundle L_λ^* (see Appendix).

The main tool to study these invariant distributions is *Bruhat’s theory*, which is presented in the Appendix. We use heavily the description of orbits of G in $S \times S \times S$ (cf. Proposition 1.2).

THEOREM 3.1. — *Let $\lambda = (\lambda_1, \lambda_2, \lambda_3)$ be three complex numbers, let $\alpha = (\alpha_1, \alpha_2, \alpha_3)$ be defined as in (2.5) and assume that they satisfy the following generic conditions*

- i) $\alpha_j \notin -\rho - 2\mathbb{N}$ for $j = 1, 2, 3$
- ii) $\alpha_1 + \alpha_2 + \alpha_3 \notin -\rho - 2\mathbb{N}$.

Then any trilinear invariant form for $\pi_{\lambda_1}, \pi_{\lambda_2}, \pi_{\lambda_3}$ is proportional to the form \mathcal{T}_λ .

Proof. — Denote by $T = T_\lambda$ the distribution density on $S \times S \times S$ for the bundle L_λ^* corresponding to an invariant trilinear form for $\pi_{\lambda_1}, \pi_{\lambda_2}, \pi_{\lambda_3}$.

Step 1: Contribution of \mathcal{O}_0

We use freely of the notation presented in the appendix. Consider the restriction T' of T to the open orbit \mathcal{O}_0 . Then \mathcal{O}_0 is a homogeneous space under G , the stabilizer H of the base point $(\mathbf{1}^+, \mathbf{1}^-, e_2)$ is compact. There is an invariant measure on G/H , the group H acts trivially on the fiber of L_λ^* , so there exists exactly one (up to constant) invariant distribution given by a smooth density. But we already know that K_α has the right

transformation property. Hence on \mathcal{O}_0 , after multiplication by a constant, we may assume that T' coincides with the restriction of \mathcal{K}_α to \mathcal{O}_0 . But the assumptions on λ guarantee that \mathcal{K}_α can be extended (by analytic continuation) as an invariant distribution on $S \times S \times S$. Hence, in order to prove that T is a multiple of \mathcal{K}_α , (i.e. to prove the uniqueness statement) we need only to prove that an invariant distribution which vanishes on \mathcal{O}_0 is identically 0. In other words, we may (and hence do) assume that

$$\text{Supp}(T) \subset \cup_{1 \leq j \leq 4} \mathcal{O}_j.$$

and proceed further to prove that $T = 0$.

Let us mention that the argument given here should be modified for the case of the circle (i.e. as $n = 2$), because there are two orbits for $SO_0(1, 2)$ in $S \times S \times S$. To restore uniqueness, one can also consider the full group $O(1, 2)$ instead. The rest of the proof is unchanged.

Step 2: Contribution of $\mathcal{O}_1, \mathcal{O}_2, \mathcal{O}_3$

We now show that

$$\text{Supp}(T) \cap \mathcal{O}_3 = \emptyset.$$

Observe that for $1 \leq j \leq 3$

$$\overline{\mathcal{O}_j} = \mathcal{O}_j \cup \mathcal{O}_4$$

so that $(\mathcal{O}_1 \cup \mathcal{O}_2 \cup \mathcal{O}_4)$ is a closed subset of $S \times S \times S$. Let

$$X = S \times S \times S \setminus (\mathcal{O}_1 \cup \mathcal{O}_2 \cup \mathcal{O}_4).$$

Then X is an open submanifold of $S \times S \times S$, acted by G , and \mathcal{O}_3 is a closed orbit of G in X . Let T'' be the restriction of T to X , so that $\text{Supp}(T'') \subset \mathcal{O}_3$. Now apply Bruhat's theory.

The normal space at the base point $(\mathbf{1}^+, \mathbf{1}^+, \mathbf{1}^-)$ is identified (via the Riemannian metric on $S \times S \times S$) with

$$N_o = \{(\xi, -\xi, 0), \xi \in T_1 S\} \simeq \mathbb{R}^{n-1}.$$

The stabilizer of the base point is $H = AM$, and there is an invariant measure on \mathcal{O}_3 , so that $\chi_0 \equiv 1$. The group M acts on N_0 by its natural action. As M modules, N_0 and its dual are equivalent, so that the spaces $S^k(N_0)$ and $\mathcal{P}_k(N_0)$ are equivalent M -modules. The space of M -invariants in $\mathcal{P}_k(N_0)$ is 0 if k is odd (and one-dimensional, generated by $|\xi|^k$ if k is even, but we won't need this fact). On the other hand, an element a_t of A acts on N_0 by dilation by e^{-t} , so that it acts on $S^k(N_0)$ by multiplication by e^{-kt} . The element a_t acts on the fiber L_0 of $L_{\chi_1, \chi_2, \chi_3}$ at $(\mathbf{1}^+, \mathbf{1}^+, \mathbf{1}^-)$

by $e^{(\rho+\lambda_1+\rho+\lambda_2-\rho-\lambda_3)t}$. Hence the element a_t acts on $S^k(N_0) \otimes L_0^*$ by multiplication by

$$e^{(-k-\rho-(\lambda_1+\lambda_2-\lambda_3))t}.$$

The assumptions on λ (more precisely $\alpha_3 = \lambda_1 + \lambda_2 - \lambda_3 \notin -\rho - 2\mathbb{N}$) guarantee that

$$\left(\mathcal{S}_k(N_0) \otimes L_0^* \otimes \mathbb{C}_{\chi_0^{-1}} \right)^H = 0$$

for any $k \in \mathbb{N}$. Hence, by Corollary A.4, there is no non-trivial invariant distribution supported in \mathcal{O}_3 .

Repeating the argument for \mathcal{O}_2 and \mathcal{O}_1 , we may (and hence do) assume now that

$$\text{Supp}(T) \subset \mathcal{O}_4.$$

Third step: Contribution of \mathcal{O}_4

Here we take $X = S \times S \times S$, as \mathcal{O}_4 is closed. The stabilizer of the base-point $(\mathbf{1}^+, \mathbf{1}^+, \mathbf{1}^+)$ is $P = MAN$. The character χ_0 is given by

$$\chi_0(ma_t n) = e^{-2\rho t}.$$

The normal space at $(\mathbf{1}^+, \mathbf{1}^+, \mathbf{1}^+)$ can be identified with

$$N_0 = \{(\xi_1, \xi_2, \xi_3), \xi_j \in T_1 S, \xi_1 + \xi_2 + \xi_3 = 0\} \simeq \mathbb{R}^{n-1} \oplus \mathbb{R}^{n-1}.$$

The group M acts on N_0 by its natural action on each factor. Again, as M modules, the space $S^k(N_0)$ is isomorphic to $\mathcal{P}_k(N_0)$. The algebra of $SO(n-1)$ invariant polynomials on $\mathbb{R}^{n-1} \oplus \mathbb{R}^{n-1}$ is generated (as an algebra) by $|\xi|^2, |\eta|^2, \langle \xi, \eta \rangle$. Hence $S^k(N_0)^M = \{0\}$ if k is odd. On the other hand, an element a_t of A acts on N_0 by multiplication by e^{-t} , hence on $S^k(N_0)$ by multiplication by e^{-kt} . It acts on the fiber L_0 by $e^{(\rho+\lambda_1+\rho+\lambda_2+\rho+\lambda_3)t}$. Hence a_t acts on $S^k(N_0) \otimes L_0^* \otimes \mathbb{C}_{\chi_0^{-1}}$ by

$$e^{(-k-\rho-(\lambda_1\lambda_2+\lambda_3))t}.$$

The assumptions on λ (namely $\lambda_1 + \lambda_2 + \lambda_3 \notin -\rho - 2\mathbb{N}$) guarantee that

$$\left(\mathcal{S}_k(N_0) \otimes L_0^* \otimes \mathbb{C}_{\chi_0^{-1}} \right)^H = \{0\}$$

for any $k \in \mathbb{N}$. Hence, by Corollary A.4, there is no non-trivial invariant distribution supported in \mathcal{O}_4 . The uniqueness statement follows. \square

4. Another construction of a trilinear invariant form

In this section we present a different construction of the invariant trilinear form \mathcal{T}_λ . Trilinear forms are connected with tensor products of representations. Roughly speaking, a trilinear form on $\mathcal{H}_1 \times \mathcal{H}_2 \times \mathcal{H}_3$ can be realized as an invariant bilinear pairing between $\mathcal{H}_1 \otimes \mathcal{H}_2$ and \mathcal{H}_3 . Although this point of view breaks the (sort of) symmetry between the three factors, it produces interesting relations with questions about tensor products of representations (cf. [12], which was the main source of inspiration for this section). Our presentation of the construction is formal and we don't work out the estimates and analytical aspects of the construction, which would follow along similar lines as in previous sections.

4.1. The space S_\mp^2 as a symmetric space

Recall that $S_\mp^2 = \{(x, y) \in S \times S, x \neq y\}$ is the orbit of $(\mathbf{1}^+, \mathbf{1}^-)$ under G . The stabilizer in G of $(\mathbf{1}^+, \mathbf{1}^-)$ is the subgroup $H = MA$.

Let $J = \begin{pmatrix} -1 & 0 & 0 & \dots & 0 \\ 0 & -1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{pmatrix}$. Then $J^t = J^{-1} = J$. The map $g \mapsto JgJ$ is an involutive automorphism of G . The set of fixed points of this involution is the subgroup $\tilde{H} = H \sqcup H^-$,

$$H^- = \left\{ h = \begin{pmatrix} \cosh t & -\sinh t & 0 \\ \sinh t & -\cosh t & 0 \\ 0 & 0 & k \end{pmatrix}, t \in \mathbb{R}, k \in O(n-1), \det k = -1 \right\}.$$

So H is the neutral component of the fixed points of an involutive automorphism of G . In other words, S_\mp^2 can be realized as the symmetric space G/H via

$$G/H \ni g \mapsto (g(\mathbf{1}^+), g(\mathbf{1}^-)).$$

There exists a unique (up to a positive real number) G -invariant measure on S_\mp^2 , namely

$$(4.1) \quad \int_{G/H} f(x) d\mu(x) = \int_S \int_S f(s, t) \frac{ds dt}{|s - t|^{2(n-1)}}.$$

Another description of G/H is as follows. Let \mathcal{X} be the set of all 2-dimensional oriented subspaces Π in $\mathbb{R}^{1,n}$ such that the restriction of q to Π is of signature $(1, 1)$. The space \mathcal{X} is an open set in the Grassmannian $G(2, n + 1)$ of oriented 2-dimensional subspaces in $\mathbb{R}^{1,n}$. The group

G operates transitively on \mathcal{X} . To the couple (s, t) in $S^2_{\mathbb{T}}$, associate the 2-dimensional space $\Pi(s, t) = \mathbb{R}\tilde{s} \oplus \mathbb{R}\tilde{t}$, with the orientation for which (\tilde{s}, \tilde{t}) is a direct basis.

PROPOSITION 4.1. — *The mapping Π is a diffeomorphism from $S^2_{\mathbb{T}}$ onto \mathcal{X} .*

Proof. — Let s, t be in $S^2_{\mathbb{T}}$. Then $\Pi(s, t)$ cannot be totally isotropic for q , because the maximally isotropic subspaces are of dimension 1. As $\Pi(s, t)$ contains two independant isotropic vectors $(\tilde{s}$ and $\tilde{t})$, the signature of q on $\Pi(s, t)$ has to be $(1, 1)$. Hence $\Pi(s, t)$ belongs to \mathcal{X} . On the other hand, let Π be in \mathcal{X} . Then the isotropic cone in Π decomposes as the union of two distinct isotropic lines, say $D \cup \Delta$. The line D (resp. Δ) intersects the hyperplane $\{x_0 = 1\}$ at a point \tilde{s} (resp. \tilde{t}) with s (resp. t) in S . Permuting D and Δ if needed, we may assume that the basis (\tilde{s}, \tilde{t}) corresponds to the orientation of Π . Then clearly $\Pi(s, t) = \Pi$. Thus Π is a 1–1 correspondance. The fact that it is a diffeomorphism is standard and left to the reader. \square

4.2. An equivariant realization of $\pi_{\sigma} \otimes \pi_{\tau}$

The group H is connected, isomorphic to $\mathbb{R} \times SO(n - 1)$. For ζ any complex number, the function ν_{ζ} on H defined by

$$\nu_{\zeta} \begin{pmatrix} \cosh t & \sinh t & 0 \\ \sinh t & \cosh t & 0 \\ 0 & 0 & k \end{pmatrix} = e^{t\zeta}$$

$(t \in \mathbb{R}, k \in M)$ is a character of H .

Form the line bundle $L_{\zeta} = G \times_H \mathbb{C}_{\zeta}$ over G/H . A section of L_{ζ} can be viewed as a C^{∞} function $F: G \rightarrow \mathbb{C}$ which satisfies

$$(4.2) \quad F(gh) = \nu_{\zeta}(h)^{-1}F(g).$$

Denote by \mathcal{L}_{ζ} the space of smooth sections of L_{ζ} over G/H . The natural action of G acts on \mathcal{L}_{ζ} by left translation is denoted by I_{ζ} :

$$(I_{\zeta}(\gamma)F)(g) = F(\gamma^{-1}g).$$

When ζ is pure imaginary, the character ν_{ζ} is unitary. If F is in \mathcal{L}_{ζ} , then $|F(gh)| = |F(g)|$ for any h in H . Hence the expression

$$(4.3) \quad |F|^2 = \int_{G/H} |F(g)|^2 d\mu(gH),$$

is well-defined (maybe $+\infty$) and is finite if (say) F has compact support modulo H . The representation I_ζ extends as a unitary representation for this inner product.

Let σ, τ be two complex numbers, and consider the representations π_σ and π_τ . As $C^\infty(S)$ is a Fréchet space, the projective and inductive topologies on the tensor product $C(S) \otimes C(S)$ coincide, and the (completed) tensor product is realized as $C^\infty(S \times S)$. Hence the tensor product $\pi_\sigma \otimes \pi_\tau$ is naturally realized on $C^\infty(S \times S)$. Explicitly,

$$(4.4) \quad \pi_\sigma \otimes \pi_\tau(g)f(x_1, x_2) = \kappa(g^{-1}, x_1)^{\rho+\sigma} \kappa(g^{-1}, x_2)^{\rho+\tau} f(g^{-1}(x_1), g^{-1}(x_2))$$

for $g \in G, f \in C^\infty(S \times S), x_1, x_2 \in S$.

For f in $C^\infty(S \times S)$, let $P_{\sigma,\tau}f$ be the function on G defined by

$$(4.5) \quad (P_{\sigma,\tau}f)(g) = \kappa(g, \mathbf{1}^+)^{\rho+\sigma} \kappa(g, \mathbf{1}^-)^{\rho+\tau} f(g(\mathbf{1}^+), g(\mathbf{1}^-)).$$

PROPOSITION 4.2. — $P_{\sigma,\tau}f$ satisfies the relation

$$(4.6) \quad (P_{\sigma,\tau}f)(gh) = \nu_{\sigma-\tau}(h)^{-1} (P_{\sigma,\tau}f)(g),$$

for g in G and h in H .

Proof. — Recall that the elements of H fix both $\mathbf{1}^+$ and $\mathbf{1}^-$. Moreover,

$$(4.7) \quad \kappa(a_t, \mathbf{1}^\pm) = e^{\mp t}, \quad t \in \mathbb{R}.$$

Now (4.6) follows (1.6). □

The map $P_{\sigma,\tau}$ will be regarded as a map from $C^\infty(S \times S)$ into $\mathcal{L}_{\sigma-\tau}$.

Let σ and τ be pure imaginary. Observe that

$$|g(\mathbf{1}^+) - g(\mathbf{1}^-)| = 2 \kappa(g, \mathbf{1}^+)^{\frac{1}{2}} \kappa(g, \mathbf{1}^-)^{\frac{1}{2}},$$

so that

$$|P_{\sigma,\tau}f|^2 = 2^{-2(n-1)} \int_{S \times S} |s - t|^{2(n-1)} |f(s, t)|^2 \frac{ds dt}{|s - t|^{2(n-1)}} = 2^{-2(n-1)} |f|^2$$

and $P_{\sigma,\tau}$ extends as an isometry (up to a constant) from $L^2(S \times S)$ onto the space of square-integrable elements of $\mathcal{L}_{\sigma-\tau}$ for the inner product associated to (4.3).

PROPOSITION 4.3. — $P_{\sigma,\tau}$ intertwines the representation $\pi_\sigma \otimes \pi_\tau$ and the representation $I_{\sigma-\tau}$, i.e. for any g in G

$$(4.8) \quad P_{\sigma,\tau} \circ (\pi_\sigma \otimes \pi_\tau)(g) = I_{\sigma-\tau}(g) \circ P_{\sigma,\tau}$$

Proof. — Let f be in $C^\infty(S \times S)$. For γ in G ,

$$P_{\sigma,\tau}((\pi_\sigma \otimes \pi_\tau)(\gamma)f)(g) = \kappa(g, \mathbf{1}^{\rho+\sigma}) \kappa(g, \mathbf{1}^-)^{\rho+\tau} \kappa(\gamma^{-1}, g(\mathbf{1}^+))^{\rho+\sigma} \dots \kappa(\gamma^{-1}, g(\mathbf{1}^-))^{\rho+\tau} f(\gamma^{-1}(g(\mathbf{1}^+), g(\mathbf{1}^-))).$$

On the other hand,

$$I_{\sigma-\tau}(\gamma)P_{\sigma,\tau}f(g) = \kappa(\gamma^{-1}g, \mathbf{1}^{\rho+\sigma}) \kappa(\gamma^{-1}g, \mathbf{1}^-)^{\rho+\tau} \dots f(\gamma^{-1}(g(\mathbf{1}^+), g^{-1}(\mathbf{1}^-))).$$

The two expressions are easily seen to be equal by using (1.6). □

4.3. Construction of a (H, ν_ζ) covariant function for π_λ

The next step consists in finding in the representation space for π_λ an element $\Theta_{\lambda,\zeta}$ which transforms under the action of H by the character ν_ζ . In general, it will be a distribution on S (a common fact in harmonic analysis on semisimple symmetric space (cf. [16])). We use a geometric approach through the realization of G/H as \mathcal{X} (see Proposition 4.1).

Let Π be an element of \mathcal{X} and s be in S . Observe that Π^\perp is a $(n-1)$ -dimensional space, complementary to Π , and the restriction of q to Π^\perp is negative-definite. Define

$$(4.9) \quad \Psi(\Pi, s) = 2(-q(\text{proj}_{\Pi^\perp} \tilde{s}))^{\frac{1}{2}}.$$

PROPOSITION 4.4. — *Let (s_1, s_2) be in S_Γ^2 . Then, for any s_3 in S ,*

$$(4.10) \quad \Psi(\Pi(s_1, s_2), s_3) = 2 \frac{|s_1 - s_3||s_2 - s_3|}{|s_1 - s_2|}.$$

Proof. — Let $\sigma_3 = \text{proj}_{\Pi^\perp} \tilde{s}_3$. Then there exist real numbers α_1 and α_2 such that $\sigma_3 = \tilde{s}_3 - \alpha_1 \tilde{s}_1 - \alpha_2 \tilde{s}_2$. The numbers α_1 and α_2 are determined by the conditions

$$[\sigma_3, \tilde{s}_1] = [\sigma_3, \tilde{s}_2] = 0.$$

Hence,

$$1 - s_3.s_2 - \alpha_1(1 - s_1.s_2) = [\tilde{s}_3, \tilde{s}_2] - \alpha_1[\tilde{s}_1, \tilde{s}_2] = [\sigma_3 + \alpha_2 \tilde{s}_2, \tilde{s}_2] = 0$$

so that

$$\alpha_1 = \frac{1 - s_2.s_3}{1 - s_1.s_2} = \left(\frac{|s_2 - s_3|}{|s_1 - s_2|} \right)^2$$

and similarly

$$\alpha_2 = \frac{1 - s_3.s_1}{1 - s_1.s_2} = \left(\frac{|s_1 - s_3|}{|s_1 - s_2|} \right)^2.$$

Now

$$\begin{aligned}
 -q(\sigma_3) &= -[\tilde{s}_3 - \alpha_1 \tilde{s}_1 - \alpha_2 \tilde{s}_2, \tilde{s}_3 - \alpha_1 \tilde{s}_1 - \alpha_2 \tilde{s}_2] \\
 &= 2\alpha_1 [\tilde{s}_1, \tilde{s}_3] + 2\alpha_2 [\tilde{s}_2, \tilde{s}_3] - 2\alpha_1 \alpha_2 [\tilde{s}_1, \tilde{s}_2] \\
 &= \left(\frac{|s_1 - s_3| |s_2 - s_3|}{|s_1 - s_2|} \right)^2.
 \end{aligned}$$

□

PROPOSITION 4.5. — *Let g be in G . Then, for any Π in \mathcal{X} and s in S ,*

$$(4.11) \quad \Psi(g(\Pi), g(s)) = \kappa(g, s)\Psi(\pi, s).$$

Proof. — As g preserves the form q and hence the orthogonality relative to q , for any Π in \mathcal{X}

$$g \circ \text{proj}_{\Pi^\perp} = \text{proj}_{g(\Pi)^\perp} \circ g.$$

Hence

$$q(\text{proj}_{g(\Pi)^\perp} g(s)) = (g\tilde{s})_0^{-2} q(\text{proj}_{g(\Pi)^\perp} g\tilde{s}) = (g\tilde{s})_0^{-2} q(\text{proj}_{\Pi^\perp} s)$$

so that

$$\Psi(g(\Pi), g(s)) = (g\tilde{s})_0^{-1} \Psi(\Pi, s) = \kappa(g, s)\Psi(\pi, s).$$

□

Let $\Pi_0 = \Pi(\mathbf{1}^+, \mathbf{1}^-) = \{\tilde{x} = (x_0, x_1, 0, \dots, 0), x_0, x_1 \in \mathbb{R}\}$, and let

$$(4.12) \quad \Psi_0(s) = \Psi(\Pi_0, s) = |\mathbf{1}^+ - s| |\mathbf{1}^- - s| = 2(x_2^2 + \dots + x_n^2)^{\frac{1}{2}}.$$

As a consequence of the previous proposition, the function Ψ_0 has a nice transformation law under the action of H .

LEMMA 4.6. — *Let $h = \begin{pmatrix} \cosh t & \sinh t & 0 \\ \sinh t & \cosh t & 0 \\ 0 & 0 & k \end{pmatrix}$, where t is in \mathbb{R} , and k is*

in $SO(n - 1)$. For any s in S

$$(4.13) \quad \Psi_0(h(s)) = \kappa(h, s)\Psi_0(s).$$

Let λ and ζ be two complex numbers. For s in S define

$$(4.14) \quad \Theta_{\lambda, \zeta}(s) = \Psi_0(s)^{-\rho - \lambda} \left| \frac{\mathbf{1}^+ - s}{\mathbf{1}^- - s} \right|^\zeta = |\mathbf{1}^+ - s|^{-\rho - \lambda + \zeta} |\mathbf{1}^- - s|^{-\rho - \lambda - \zeta}.$$

PROPOSITION 4.7. — *The function $\Theta_{\lambda, \zeta}$ satisfies*

$$(4.15) \quad \pi_\lambda(h)\Theta_{\lambda, \zeta} = \nu_\zeta(h)\Theta_{\lambda, \zeta}$$

for all h in H .

Proof. — Let h be in H . Then

$$\begin{aligned} \pi_\lambda(h)\Theta_{\lambda,\zeta}(s) &= \kappa(h^{-1}, s)^{\rho+\lambda} \Theta_{\lambda,\zeta}(h^{-1}(s)) \\ &= \kappa(h^{-1}, s)^{\rho+\lambda} \Psi_0(h^{-1}(s))^{-\rho-\lambda} \left| \frac{\mathbf{1}^+ - h^{-1}(s)}{\mathbf{1}^- - h^{-1}(s)} \right|^\zeta \\ &= \kappa(h^{-1}, s)^{\rho+\lambda} \kappa(h^{-1}, s)^{-\rho-\lambda} \Psi_0(s)^{-\rho-\lambda} \left| \frac{\kappa(h^{-1}, \mathbf{1}^+)}{\kappa(h^{-1}, \mathbf{1}^-)} \right|^{\frac{\zeta}{2}} \left| \frac{\mathbf{1}^+ - s}{\mathbf{1}^- - s} \right|^\zeta \\ &= \Psi_0(s)^{-\rho-\lambda} (e^{2t})^{\frac{\zeta}{2}} \left| \frac{\mathbf{1}^+ - s}{\mathbf{1}^- - s} \right|^\zeta = \nu_\zeta(h) \Theta_{\lambda,\zeta}(s) \end{aligned}$$

by using (4.7). □

4.4. The duality between I_ζ and π_λ

Now define the corresponding *Fourier transform* $\mathcal{F}_{\lambda,\zeta}$: For F a smooth section of L_ζ with compact support modulo H , define $\mathcal{F}_{\lambda,\zeta}F$ by

$$(4.16) \quad \mathcal{F}_{\lambda,\zeta}F(s) = \int_{G/H} F(g) \pi_\lambda(g) \Theta_{\lambda,\zeta}(s) d\mu(gH).$$

Observe that, thanks to (4.15) the integrand is a function on G/H , so that the integral makes sense, the result being in general a distribution on S .

PROPOSITION 4.8. — *For any g in G*

$$(4.17) \quad \mathcal{F}_{\lambda,\zeta} I_\zeta(g) = \pi_\lambda(g) \mathcal{F}_{\lambda,\zeta} F.$$

Proof. — Set $\Theta = \Theta_{\lambda,\zeta}$, $\mathcal{F} = \mathcal{F}_{\lambda,\zeta}$, let F be in \mathcal{L}_ζ with compact support modulo H and let γ be in G . Then

$$\mathcal{F} \circ I_\zeta(\gamma)F(s) = \int_{G/H} F(\gamma^{-1}g) \kappa(g^{-1}, s)^{\rho+\lambda} \Theta(g^{-1}(s)) d\mu(gH).$$

Set $\gamma^{-1}g = l$ and use the invariance of the measure $d\mu$ to obtain

$$\begin{aligned} \mathcal{F} \circ I_\zeta(\gamma)F(s) &= \int_{G/H} F(l) \kappa(l^{-1}\gamma^{-1}, s)^{\rho+\lambda} \Theta(l^{-1}(\gamma^{-1}(s))) d\mu(lH) \\ &= \kappa(\gamma^{-1}, s)^{\rho+\lambda} \int_{G/H} F(l) \kappa(l^{-1}, \gamma^{-1}(s))^{\rho+\lambda} \Theta(l^{-1}(\gamma^{-1}(s))) d\mu(lH) \\ &= \kappa(\gamma^{-1}, s)^{\rho+\lambda} \int_{G/H} F(l) \pi_\lambda(l) \Theta(\gamma^{-1}(s)) d\mu(lH) \\ &= \kappa(\gamma^{-1}, s)^{\rho+\lambda} \mathcal{F}F(\gamma^{-1}(s)) = (\pi_\lambda(\gamma)\mathcal{F}F)(s). \end{aligned}$$

□

4.5. Application to trilinear forms

THEOREM 4.9. — *Let $\lambda_1, \lambda_2, \lambda_3$ be three complex numbers. For f_1, f_2, f_3 three functions in $\mathcal{C}^\infty(S)$, let $T(f_1, f_2, f_3)$ be defined by*

$$(4.18) \quad T(f_1, f_2, f_3) = \left(\mathcal{F}_{-\lambda_3, \lambda_1 - \lambda_2} P_{\lambda_1, \lambda_2}(f_1 \otimes f_2), f_3 \right).$$

Then, for any $g \in G$,

$$T(\pi_{\lambda_1}(g)f_1, \pi_{\lambda_2}(g)f_2, \pi_{\lambda_3}(g)f_3) = T(f_1, f_2, f_3),$$

whenever the right hand side is well defined.

Proof. — Let g be in G and let f_1, f_2, f_3 be three functions in $\mathcal{C}^\infty(S)$. Then

$$\begin{aligned} & T(\pi_{\lambda_1}(g)f_1, \pi_{\lambda_2}(g)f_2, \pi_{\lambda_3}(g)f_3) \\ &= \left(\mathcal{F}_{-\lambda_3, \lambda_2 - \lambda_1}, P_{\lambda_1, \lambda_2}(\pi_{\lambda_1}(g)f_1 \otimes \pi_{\lambda_2}(g)f_2), \pi_{\lambda_3}(g)f_3 \right) \\ &= \left(\mathcal{F}_{-\lambda_3, \lambda_1 - \lambda_2} I_{\lambda_1 - \lambda_2}(g) P_{\lambda_1, \lambda_2}(f_1 \otimes f_2), \pi_{\lambda_3}(g)f_3 \right) \text{ (use (4.8))} \\ &= \left(\pi_{-\lambda_3}(g) (\mathcal{F}_{-\lambda_3, \lambda_1 - \lambda_2} P_{\lambda_1, \lambda_2}(f_1 \otimes f_2)), \pi_{\lambda_3}(g)f_3 \right) \text{ (use (4.17))} \\ &= T(f_1, f_2, f_3) \text{ (use (1.13)).} \end{aligned}$$

□

Making explicit the right handside of (4.18) shows that it coincides (up to a constant) with the former expression of \mathcal{T}_λ where $\lambda = (\lambda_1, \lambda_2, \lambda_3)$ (see Theorem 2.1).

Appendix A. Invariant distributions supported by a submanifold

Here is a presentation of the main results in Bruhat’s theory [2], but written in terms of vector bundles and distribution densities (in the sense of [7] ch. VI). We sketch the main steps of the proof, following [18]. See also [8] for more results on this topic.

A.1. Distribution densities for a vector bundle over a manifold

First recall the composition of a distribution with a diffeomorphism. Let X_1 and X_2 be two open sets of \mathbb{R}^N and let $\Phi: X_1 \rightarrow X_2$ be a \mathcal{C}^∞ diffeomorphism. Then there is a unique *continuous* linear map $\Phi^*: \mathcal{D}'(X_2) \rightarrow \mathcal{D}'(X_1)$

which extends the composition of functions, i.e. such that $\Phi^*f = f \circ \Phi$ for $f \in \mathcal{C}(X_2)$.

A *distribution density* on a manifold X is by definition a continuous linear form on $\mathcal{C}_c^\infty(X)$. Let u be a distribution density on X . Let (X_κ, κ) be a local chart, i.e. κ is a diffeomorphism of an open set X_κ of X onto an open set \tilde{X}_κ of \mathbb{R}^N . Then the formula

$$(A.1) \quad u_\kappa(\varphi) = u(\varphi \circ \kappa)$$

for $\varphi \in \mathcal{C}_c^\infty(\tilde{X}_\kappa)$ defines a distribution u_κ on \tilde{X}_κ , called the *local expression* of u in the chart (X_κ, κ) . Further, let (X_κ, κ) and $(X_{\kappa'}, \kappa')$ be two overlapping charts, and let u_κ and $u_{\kappa'}$ the corresponding local expressions of u . Let

$$\Phi: \kappa'(X_\kappa \cap X_{\kappa'}) \rightarrow \kappa(X_\kappa \cap X_{\kappa'})$$

be the change of coordinates (equal to $\kappa \circ \kappa'^{-1}$). Then

$$(A.2) \quad u_{\kappa'} = |\det d\Phi| \Phi^* u_\kappa \quad \text{in } \kappa'(X_\kappa \cap X_{\kappa'}).$$

Conversely, suppose we have an atlas \mathcal{F} of charts (X_κ, κ) covering the manifold M and suppose that for each κ we are given a distribution $u_\kappa \in \mathcal{D}'(\tilde{X}_\kappa)$. Assume further that for any two overlapping charts (X_κ, κ) and $(X_{\kappa'}, \kappa')$, the condition (A.2) is satisfied. Then the system $((u_\kappa), \kappa \in \mathcal{F})$ defines a unique distribution density u on X such that, for κ in \mathcal{F} and $\varphi \in \mathcal{C}_c^\infty(\tilde{X}_\kappa)$ condition (A.1) is satisfied.

The space of distribution densities on X is denoted by $\mathcal{D}'(X)$. A *smooth density* is a density the local expressions of which are \mathcal{C}^∞ functions. The smooth densities are \mathcal{C}^∞ sections of a line bundle called the *density bundle* $\Omega(X)$. It is very similar to the bundle of differential forms of maximal degree on X , in the sense that their transition functions just differ by an absolute value.

This definition can be extended to the case of vector bundles. Let $\pi: L \rightarrow X$ be a \mathcal{C}^∞ vector bundle over M , with model fiber E_0 . Let (X_i) be a family of open subsets of X such that over each X_i the bundle can be trivialized. Let $\Psi_i: \pi^{-1}(X_i) \rightarrow X_i \times E_0$ and $\Psi_j: \pi^{-1}(X_j) \rightarrow X_j \times E_0$ be two trivializations of the bundle over two overlapping subsets X_i and X_j . Then the map $\Psi_i \circ \Psi_j^{-1}$ over $(X_i \cap X_j) \times L_0 \rightarrow (X_i \cap X_j) \times L_0$ is of the form

$$(x, v) \mapsto (x, g_{ij}(x)v)$$

where $g_{ij}(x)$ is in $GL(L_0)$ and the map (*transition functions of the bundle*)

$$g_{ij}: X_i \cap X_j \ni x \mapsto g_{ij}(x) \in GL(E_0)$$

is \mathcal{C}^∞ . A distribution density for the bundle L is a system (u_i) of distribution densities on X_i with values in L_0 such that

$$u_i = g_{ij}u_j \quad \text{in } X_i \cap X_j.$$

Denote by $\mathcal{D}'(X, L)$ the space of distribution densities for the bundle L .

Let \mathcal{L}_c^∞ be the space of C^∞ sections with compact support of the bundle L . Then the dual of \mathcal{L}_c^∞ is identified with the space $\mathcal{D}'(X, L^*)$. The smooth elements in the dual (those given locally by integration against a smooth function) are the \mathcal{C}^∞ sections of the bundle $\Omega(X) \otimes L^*$.

A.2. Invariant distribution: The case of a homogeneous vector bundle

Let G be a Lie group acting transitively on a manifold X . Let o be a base-point in X , and $H = G^o$ be its stabilizer in G , so that $X \simeq G/H$. An element h of H acts on X and fixes o , so that by differentiation, it acts on its tangent plane by (say) $\tau_0(h)$. The tangent space T_0X can be identified with $\mathfrak{g}/\mathfrak{h}$. The element h acts on \mathfrak{g} by the adjoint action $\text{Ad}_{\mathfrak{g}}(h)$, and preserves the subalgebra \mathfrak{h} on which it acts by $\text{Ad}_{\mathfrak{h}} h$. Hence it acts on $\mathfrak{g}/\mathfrak{h}$, and this action coincides with $\tau_0(h)$. This action satisfies

$$(A.3) \quad \det \tau_0(h) = \frac{\det(\text{Ad}_{\mathfrak{g}} h)}{\det(\text{Ad}_{\mathfrak{h}} h)}.$$

The modular function δ_G of a Lie group G is defined by

$$\delta_G(g) = |\det \text{Ad}(g^{-1})|$$

so that

$$(A.4) \quad |\det \tau_0(h)| := \chi_0(h) = \frac{\delta_H(h)}{\delta_G(h)}.$$

A homogeneous vector bundle L over X is a vector bundle L together with an action of the group G on L by bundle isomorphisms. If g is in G and x in X , then g maps the fiber L_x into $L_{g(x)}$ by a linear isomorphism. In particular, H acts on L_o by a representation (say) τ . Conversely, given a representation τ of H in a vector space E , then one constructs the bundle $G \times_\tau E$ as $G \times E / \sim$, where \sim is the equivalence relation defined by the right action of H on $G \times E$

$$(g, v) \sim (gh^{-1}, \tau(h)v) \quad \text{for some } h \in H,$$

g in G and $v \in E$. Any homogeneous vector bundle over X is of that sort, in the sense that the bundle L is isomorphic to $G \times_\tau L_0$ (see [17]).

A section $s: X \rightarrow L$ can be realized as a L_0 -valued function f_s on G which satisfies

$$(A.5) \quad f_s(gh) = \tau^{-1}(h)f_s(g)$$

and, conversely, such a function f gives rise to a section of L . Let \mathcal{L}_c^∞ be the space of smooth sections of L with compact support. The space \mathcal{L}_c^∞ is G -equivariantly isomorphic to $\mathcal{C}_c^\infty(G, H, \tau)$, the space of \mathcal{C}^∞ functions on G which satisfy (A.5) and have compact support modulo H . The group G acts by left translations on \mathcal{L}_c^∞ , and this action is equivariant with the left action of G on $\mathcal{C}_c^\infty(G, H, \tau)$.

The tangent bundle TX of X is an example of such a homogeneous bundle. The action of H on the fiber T_0X is τ_0 . Another important homogeneous bundle is the bundle $\Omega(X)$ of densities over X . It is a line bundle, corresponding to the character of H given by $|\det((\tau(h)^{-1})^t)| = \chi_o(h)^{-1}$ (cf. (A.4)), so that, in this context, we denote the fiber at o of the bundle $\Omega(X)$ by $\mathbb{C}_{\chi_o^{-1}}$.

THEOREM A.1. — *There exists a non trivial invariant continuous linear forms on \mathcal{L}_c^∞ if and only if there exists a non trivial linear form ξ on E such that, for all $h \in H$*

$$(A.6) \quad \tau(h^{-1})^t \xi = \chi_o(h)\xi.$$

More precisely,

$$(A.7) \quad (\mathcal{L}_c^{\infty*})^G = \mathcal{D}'(X, L^*)^G \simeq (L_0^* \otimes \mathbb{C}_{\chi_o^{-1}})^H.$$

Sketch of proof. — An element of the dual of \mathcal{L}_c^∞ is a distribution density for the bundle L^* . If it is invariant by G , then the corresponding distribution density turns out to be smooth. Hence we are looking for a G -invariant smooth section of the bundle $L^* \otimes \Omega(X)$. But this is equivalent to an H invariant element in the fiber at o . □

A.3. Invariant distribution supported in a submanifold

Let X be a manifold and G a Lie group acting on X . Let Q be an orbit of G in X and assume that Q is closed. Let L be a homogeneous vector bundle over X . Let N be the normal bundle of Q (i.e. the quotient bundle $TX|_Q/TQ$). Fix a base-point o in Q , let $G_o = H$ be the stabilizer of o . Then H acts on the tangent space T_oX , preserving the subspace T_oQ , and hence acts on the normal space $N_o = T_oX/T_oQ$ at o .

Let T be a distribution on Q . Then the map

$$\mathcal{L}_c^\infty(X) \ni \varphi \longmapsto (T, \varphi|_Q)$$

defines a distribution on X , which we still denote by T .

Let \mathcal{L}_c^∞ be the space of smooth sections of L with compact support. Let T be a continuous linear form on \mathcal{L}_c^∞ supported in Q . Choose a local coordinate system on X

$$(u_1, \dots, u_s, v_1, \dots, v_r)$$

such that $v_j = 0, 1 \leq j \leq r$ are local equations for Q . The $(u_i)_{1 \leq i \leq s}$ form a coordinate system of Q near o . On the other hand let N be the normal bundle of Q . The family $(\frac{\partial}{\partial v_j})_{1 \leq j \leq r}$ gives a local trivialization of the normal bundle N . Choose a local trivialization of the bundle L , and denote by e_1^*, \dots, e_l^* the corresponding coordinates on the fiber.

By Schwarz's local structure theorem for distributions supported in a vector subspace, there exists an integer $k \in \mathbb{N}$ (the local transversal order of the distribution), and for each $j, 1 \leq j \leq l$ and each multiindex $\alpha = (\alpha_1, \dots, \alpha_r)$ with $|\alpha| = \alpha_1 + \dots + \alpha_r \leq k$ uniquely determined distributions T_α^j on (some open subset of) $\mathbb{R}^s \subset \mathbb{R}^s \times \mathbb{R}^r$ such that

$$T = \sum_{1 \leq j \leq l} \sum_{|\alpha| \leq k} (-1)^{|\alpha|} D^\alpha T_\alpha^j e_j^*$$

Let us consider the "top terms" subcollection $(T_\alpha^j, |\alpha| = k, 1 \leq j \leq l)$. This collection can be interpreted as the local expression of a distribution density for the bundle $S^k(N) \otimes L_{|Q}^*$ on Q . This is obtained by checking the way the collection transforms under

- change of local trivialization of the bundle L (change the e_j^* 's)
- change of local trivialization of the normal bundle (change the v_j 's)
- change of the local coordinate system on Q (change the u_j 's).

Denote by $\sigma^{(k)}(T)$ the section of the bundle $S^k(N) \otimes L_{|Q}^*$ associated to T .

Denote by $\mathcal{D}'_k(Q, L)$ the space of continuous linear forms on \mathcal{L}_c^∞ which are supported on Q and of transversal order $\leq k$ on any local chart of X .

THEOREM A.2. — *The map $T \longmapsto \sigma^{(k)}(T)$ is a linear map from $\mathcal{D}'_k(Q, L)$ in $\mathcal{D}'(Q, S^k(N) \otimes L_{|Q}^*)$, with kernel $\mathcal{D}'_{k-1}(Q, L)$. Moreover, if Φ is a bundle diffeomorphism of the bundle L , which maps Q into Q , then*

$$\sigma^{(k)}(\Phi^*T) = \Phi^* \sigma^{(k)}(T),$$

where Φ^* denotes the action naturally induced by Φ on $\mathcal{D}'(Q, L)$ or of the restriction of $\Phi|_{L|Q}$ on $\mathcal{D}'(Q, S^k(N) \otimes L_{|Q}^*)$.

Assume now that T is invariant by G . Observe that invariance insures that the local transversal degree of T is the same for all charts of Q .

Denote by $\mathcal{D}'_k(Q, L)^G$ the space of G -invariant elements of $\mathcal{D}'_k(Q, L)$. The conjunction of Theorem A.1 and Theorem A.2 gives some estimate of its dimension.

THEOREM A.3. — For any $k \in \mathbb{N}$,

$$\dim(\mathcal{D}'_k(Q, L)^G / \mathcal{D}'_{k-1}(Q, L)^G) \leq \dim(\mathcal{S}_k(N_0) \otimes L_0^* \otimes \mathbb{C}_{\chi_0^{-1}})^H.$$

The theorem is mostly used through the following corollary.

COROLLARY A.4. — Assume that for any $k \in \mathbb{N}$

$$(\mathcal{S}_k(N_0) \otimes L_0^* \otimes \mathbb{C}_{\chi_0^{-1}})^H = \{0\}.$$

Then there exists no non trivial continuous G -invariant linear form on \mathcal{L}_c^∞ such that $\text{Supp}(T) \subset Q$.

BIBLIOGRAPHY

- [1] J. BERNSTEIN & A. REZNIKOV, “Estimates of automorphic functions”, *Mosc. Math. J.* **4** (2004), no. 1, p. 19-37.
- [2] F. BRUHAT, “Sur les représentations induites des groupes de Lie”, *Bull. Soc. Math. France* **84** (1956), p. 97-205.
- [3] J.-L. CLERC, T. KOBAYASHI, B. ØRSTED & M. PEVZNER, “Generalized Bernstein-Reznikov integrals”, to be published in *Mathematische Annalen*, DOI 10.1007/s0028-010-0516-4.
- [4] J.-L. CLERC & K.-H. NEEB, “Orbits of triples in the Shilov boundary of a bounded symmetric domain”, *Transform. Groups* **11** (2006), no. 3, p. 387-426.
- [5] A. DEITMAR, “Invariant triple products”, *Int. J. Math. Math. Sci.* (2006), p. 22, Art. ID 48274.
- [6] I. M. GEL'FAND & G. E. SHILOV, *Generalized functions. Vol. 1*, Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1964 [1977], Properties and operations, Translated from the Russian by Eugene Saletan, xviii+423 pages.
- [7] L. HÖRMANDER, *The analysis of linear partial differential operators. I*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 256, Springer-Verlag, Berlin, 1983, Distribution theory and Fourier analysis, ix+391 pages.
- [8] J. A. C. KOLK & V. S. VARADARAJAN, “On the transverse symbol of vectorial distributions and some applications to harmonic analysis”, *Indag. Math. (N.S.)* **7** (1996), no. 1, p. 67-96.
- [9] P. LITTELMANN, “On spherical double cones”, *J. Algebra* **166** (1994), no. 1, p. 142-157.
- [10] H. Y. LOKE, “Trilinear forms of \mathfrak{gl}_2 ”, *Pacific J. Math.* **197** (2001), no. 1, p. 119-144.
- [11] P. MAGYAR, J. WEYMAN & A. ZELEVINSKY, “Multiple flag varieties of finite type”, *Adv. Math.* **141** (1999), no. 1, p. 97-118.
- [12] V. F. MOLČANOV, “Tensor products of unitary representations of the three-dimensional Lorentz group”, *Izv. Akad. Nauk SSSR Ser. Mat.* **43** (1979), no. 4, p. 860-891, 967.

- [13] A. I. OKSAK, “Trilinear Lorentz invariant forms”, *Comm. Math. Phys.* **29** (1973), p. 189-217.
- [14] C. SABBABH, “Polynômes de Bernstein-Sato à plusieurs variables”, in *Séminaire sur les équations aux dérivées partielles 1986–1987*, École Polytech., Palaiseau, 1987, Exp. No. XIX, 6.
- [15] R. TAKAHASHI, “Sur les représentations unitaires des groupes de Lorentz généralisés”, *Bull. Soc. Math. France* **91** (1963), p. 289-433.
- [16] E. P. VAN DEN BAN, “The principal series for a reductive symmetric space. I. H -fixed distribution vectors”, *Ann. Sci. École Norm. Sup. (4)* **21** (1988), no. 3, p. 359-412.
- [17] N. R. WALLACH, *Harmonic analysis on homogeneous spaces*, Marcel Dekker Inc., New York, 1973, Pure and Applied Mathematics, No. 19, xv+361 pages.
- [18] G. WARNER, *Harmonic analysis on semi-simple Lie groups. I*, Springer-Verlag, New York, 1972, Die Grundlehren der mathematischen Wissenschaften, Band 188, xvi+529 pages.

Manuscrit reçu le 1^{er} mars 2010,
accepté le 18 mai 2010.

Jean-Louis CLERC
Université Henri Poincaré (Nancy 1)
Institut Élie Cartan
54506 Vandoeuvre-lès-Nancy (France)
jlclerc@iecn.u-nancy.fr
Bent ØRSTED
Matematisk Institut
Byg. 430, Ny Munkegade
8000 Aarhus C (Denmark)
orsted@imf.au.dk