



ANNALES

DE

L'INSTITUT FOURIER

Bernard SHIFFMAN, Tatsuya TATE & Steve ZELDITCH

Distribution laws for integrable eigenfunctions

Tome 54, n° 5 (2004), p. 1497-1546.

http://aif.cedram.org/item?id=AIF_2004__54_5_1497_0

© Association des Annales de l'institut Fourier, 2004, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

DISTRIBUTION LAWS FOR INTEGRABLE EIGENFUNCTIONS

by Bernard SHIFFMAN, Tatsuya TATE
and Steve ZELDITCH

Contents.

1. Introduction	1498
2. Background on toric varieties and moment polytopes	1506
2.1. The line bundle L_p^c and associated circle bundle X_P^c	1508
2.2. Moment maps and torus actions	1510
2.3. Fourier analysis	1511
2.4. Monomials on projective space	1513
3. Pointwise asymptotics on general toric varieties	1517
3.1. Pointwise asymptotics: interior points	1517
3.2. Pointwise asymptotics: boundary lattice points	1522
3.3. Moments and \mathcal{L}^{2k} norms	1529
3.4. Asymptotics on projective space	1534
4. Asymptotics of distribution functions	1536
4.1. Rescaled distributions functions	1537
4.2. Non-rescaled distribution functions	1540
Bibliography	1545

Research partially supported by NSF grants DMS-0100474 (first author) and DMS-0071358 (third author) and by JSPS (second author).

Keywords: Toric Kähler variety – Joint eigenfunction of the torus action – Distribution law of eigenfunctions – Semi-classical scaling limit.

Math. classification: 35P20 – 32H30 – 81S30.

1. Introduction.

A problem of considerable interest in both mathematics and physics is to determine the asymptotics of the distribution functions

$$D_j(t) := \text{Vol}\{z : |\varphi_j(z)|^2 > t\}$$

of an orthonormal basis $\{\varphi_j\}$ of eigenfunctions of a Laplacian (or similar Hamiltonian) on a compact manifold (see [Y]). In general, it is hopelessly difficult to obtain more than crude bounds on such distribution functions, which of course control the \mathcal{L}^p -norms of the eigenfunctions. Numerical analyses and heuristics from quantum chaos and disordered systems suggest however a rich picture in which the asymptotics of $D_j(t)$ is related to the classical dynamics underlying the eigenvalue problem. (Some references will be discussed at the end of the introduction.) The purpose of this paper is to give a rather complete analysis of the limit distribution of eigenfunctions in one of the few settings where such a detailed analysis is possible, namely where the phase space is a toric Kähler variety (M, ω) .

Let us recall the definitions (see §2 for details). Toric varieties are complex manifolds on which the complex torus $(\mathbb{C}^*)^m$ acts with an open dense orbit. By a toric Kähler variety we mean a toric variety equipped with a Kähler form (M, ω) that is invariant under the underlying real torus \mathbf{T}^m . The action of \mathbf{T}^m is Hamiltonian with respect to ω , and thus toric varieties are models of completely integrable systems. They are of a very special type because integrable systems usually generate an \mathbb{R}^m action rather than a \mathbf{T}^m action. Although there are rigidity theorems limiting the class of such examples in the world of real Riemannian manifolds [LS], toric varieties provide a plentiful collection in the world of complex manifolds.

The torus action can be ‘quantized’ or linearized on the Hilbert space completion of the coordinate ring

$$(1) \quad \mathcal{H} := \bigoplus_{N=0}^{\infty} H^0(M, L^N),$$

where $L \rightarrow M$ is a holomorphic line bundle with $c_1(L) = \frac{1}{2\pi} \omega$ and where $H^0(M, L^N)$ denotes the space of holomorphic sections of its N -th tensor power. This quantization is generally known as the holomorphic (Bargmann-Fock) representation in the physics literature. The space \mathcal{H} is spanned by joint eigenfunctions of the linearized $(\mathbb{C}^*)^m$ action, which we

refer to as ‘monomials.’ In the fundamental case of $M = \mathbb{C}P^m$, the joint eigenfunctions are the monomials given in an affine chart by

$$(2) \quad \chi_\alpha : \mathbb{C}^m \rightarrow \mathbb{C}, \quad \chi_\alpha(z) = z^\alpha.$$

The monomials lift (by homogenization) to homogeneous monomials on \mathbb{C}^{m+1} .

We consider the case where M is a smooth projective toric variety; i.e., $M = M_P, L = L_P$, where P is an integral Delzant polytope (see §2). Then the linearized \mathbf{T}^m action is generated by m commuting operators $\hat{I}_j, j = 1, \dots, m$ on M_P which preserve $H^0(M_P, L_P^N)$, and the joint spectrum of the eigenvalue problem

$$(3) \quad \hat{I}_j \varphi_\alpha^P = \alpha_j \varphi_\alpha^P, \quad \alpha = (\alpha_1, \dots, \alpha_m) \in \mathbb{R}^m, \quad \varphi_\alpha^P \in H^0(M_P, L_P^N)$$

consists of lattice points $\alpha \in NP \cap \mathbb{Z}^m$. Here and below, φ_α^P denotes the \mathcal{L}^2 -normalized joint eigenfunction.

Our main results concern the asymptotics of their distribution functions

$$(4) \quad D_\gamma(t) := \text{Vol}\{z \in M_P : |\varphi_\gamma^P(z)|^2 > t\}$$

with $\gamma \in NP \cap \mathbb{Z}^m$ as $N \rightarrow \infty$. The function $|\varphi_\gamma^P(z)|^2$ is often called the ‘Husimi distribution’ in the physics literature, and thus our results determine its distribution law. The norm $|\varphi_\gamma^P(z)|$ of $\varphi_\gamma^P(z) \in L_P^N$ is the pull-back of the Fubini-Study norm under a monomial embedding of the form

$$(5) \quad \begin{aligned} \Phi_P^c &= [c_{\alpha(1)}\chi_{\alpha(1)}, \dots, c_{\alpha(d+1)}\chi_{\alpha(d+1)}] : (\mathbb{C}^*)^m \rightarrow \mathbb{C}P^d, \\ P \cap \mathbb{Z}^m &= \{\alpha(1), \dots, \alpha(d+1)\}, \end{aligned}$$

for a choice of constants $c_{\alpha(j)} \in \mathbb{C}^*$. The volume in M_P and the Hermitian norm h_P^c on L_P are by definition the pull-backs of the Fubini-Study metric and form under this monomial embedding, and the \mathcal{L}^2 norm on the space $H^0(M, L_P^N)$ is in turn induced from the volume form and the Hermitian pointwise norm of h_P^c . (See §2 for details.)

As Figure 1 illustrates, the monomial φ_α^P is something like a Gaussian bump centered on the real torus $\mu_P^{-1}(\alpha)$, where $\mu_P : M_P \rightarrow P$ is the moment map for the classical Hamiltonian \mathbf{T}^m -action on M_P (see §1).




Figure 1. $2\pi/N$ times the monomial $|\varphi_{N\alpha}^{\Sigma}(z)|^2$ for $N = 1$ (left) and $N = \infty$ (right) for $m = 2$ and $\alpha = (2, 3)$, where Σ is the standard simplex. The variable z is chosen as $u \mapsto z = e^{(\rho_\alpha + u/\sqrt{N})/2}$. See Proposition 3.17 in Section 3.

Since we would like to determine the properties of eigenfunctions $\varphi_\alpha \in NP$ when α is large, but not necessarily a multiple of a lattice point in P , we shall consider a sequence of approximate multiples, as in the following definition:

DEFINITION 1.1. — Let $\alpha_N \in NP \cap \mathbb{Z}^m$ be a sequence of lattice points, and let $x \in P$. We say that $\{\alpha_N\}$ is a sequence of approximate multiples of x if

$$(6) \quad \alpha_N = Nx + O(1).$$

Our first result gives the pointwise behavior of the eigenfunctions:

THEOREM 1.2. — Let x be a point in the interior P° of the polytope P , which is not necessarily a lattice point. Then there exists a non-negative function $b_x^P \in C^\infty((\mathbb{C}^*)^m)$ such that $b_x^P(z) = 0$ if and only if $z \in \mu_P^{-1}(x)$,

and for every sequence $\alpha_N \in NP \cap \mathbb{Z}^m$ of approximate multiples of x , we have

$$|\varphi_{\alpha_N}^P(z)|^2 = c(P, x) \left(\frac{N}{2\pi}\right)^{m/2} e^{-N(b_x^P(z) - \langle \tau_x^P(z), x_N \rangle)} [1 + O(N^{-1})]$$

uniformly on $(\mathbb{C}^*)^m$, where $c(P, x) \in \mathbb{R}^+$ and $x_N = \alpha_N/N - x$.

The constant $c(P, x)$ is defined in (43) and (45) (and also appears in Theorems 1.3 and 1.4).

When we consider a sequence of the lattice points of the form $\alpha_N = N\alpha$ with a lattice point α , the assumption that $\alpha \in P^\circ$ is not necessary. In fact, we will give the pointwise asymptotics for $\alpha_N = N\alpha$ with any lattice point $\alpha \in P \cap \mathbb{Z}^m$ in Section 3 (see Propositions 3.5 and 3.8).

In the above Theorem, the function $b_x^P(z)$ ($x \in P^\circ$) on $(\mathbb{C}^*)^m$ is defined by

$$(7) \quad b_x^P(z) := \log \left(\frac{\sum_{\beta \in P} |c_\beta z^\beta|^2}{\sum_{\beta \in P} e^{-\langle \tau_x^P(z), \beta \rangle} |c_\beta z^\beta|^2} \right) - \langle \tau_x^P(z), x \rangle,$$

where $\tau_x^P(z) \in \mathbb{R}^m$ is the vector given by the equation

$$(8) \quad \mu_P(e^{-\tau_x^P(z)/2} z) = x, \quad z \in (\mathbb{C}^*)^m.$$

Here μ_P is the moment map for the \mathbf{T}^m action on M_P (see (23) for the definition), and we write

$$(9) \quad e^r z = (e^{r_1} z_1, \dots, e^{r_m} z_m) \quad \text{for } r \in \mathbb{R}^m, z \in (\mathbb{C}^*)^m.$$

We can express (as in [SZ2(17)–(18)]) the function b_x^P in the more intuitive form as follows: we introduce the real power ‘monomials’

$$|\chi_x(z)| := |z|^x = |z_1|^{x_1} \dots |z_m|^{x_m}$$

and define

$$(10) \quad \mathcal{M}_x^P(z) := \frac{|\chi_x(z)|_P}{\sup |\chi_x(z)|_P},$$

where (cf. (20))

$$|\chi_x(z)|_P := \frac{|\chi_x(z)|}{\sqrt{\sum_{\beta \in P} |c_\beta z^\beta|^2}} \quad (z \in (\mathbb{C}^*)^m).$$

(The normalized monomial \mathcal{M}_x^P has sup-norm 1, attained on the torus $\mu_P^{-1}(x)$.) Then (7) is equivalent to:

$$(11) \quad b_x^P(z) = -2 \log \mathcal{M}_x^P(z).$$

As $N \rightarrow \infty$, the sequence of \mathcal{L}^2 -normalized monomials $\varphi_{N\alpha}^P$ flattens out exponentially quickly away from the peak set $\mu_P^{-1}(\alpha)$. Hence their distribution functions tend to zero as $N \rightarrow \infty$ for any fixed $t > 0$. The rate of decay of $D_{N\alpha}(t)$ as $N \rightarrow \infty$ is given by the following result:

THEOREM 1.3. —

(i) Let $\alpha_N \in NP \cap \mathbb{Z}^m$ be a sequence of lattice points which are approximate multiples of $x \in P^\circ$ (see (6)). Then, for $t > 0$, we have

$$D_{\alpha_N}(t) \sim \frac{(\pi m)^{m/2}}{c(P, x)\Gamma(m/2 + 1)} \left(\frac{\log N}{N}\right)^{m/2}.$$

(ii) Let $\alpha \in \partial P \cap \mathbb{Z}^m$. Then, for $t > 0$, we have

$$D_{N\alpha}(t) \sim \frac{(\pi d(\alpha))^{d(\alpha)/2}}{c(P, \alpha)\Gamma(d(\alpha)/2 + 1)} \left(\frac{\log N}{N}\right)^{d(\alpha)/2},$$

where $d(\alpha) := m + \text{codim } F_\alpha$, F_α being the face of P containing α .

Here \sim means the ratio of the left and right hand sides tends to 1, and the constants $c(P, x)$, $c(P, \alpha)$ are given by (43) and (73)–(45). We recall that if x is a point in a face F of codimension r , then $\mu_P^{-1}(x) \cong \mathbf{T}^{m-r}$. Hence $d(\alpha) = 2m - \dim \mu_P^{-1}(\alpha)$.

The exponentially localized behavior of the monomials suggests studying the distribution function on various length scales. First, we show that the D_{α_N} have a universal scaling limit on a small length scale:

THEOREM 1.4. —

(i) Let $\alpha_N \in NP \cap \mathbb{Z}^m$ be a sequence of lattice points satisfying the condition (6) with some point $x \in P^\circ$. Then, for $0 < t \leq c(P, x)$, we have

$$\lim_{N \rightarrow \infty} (N/2\pi)^{m/2} D_{\alpha_N} \left((N/2\pi)^{m/2} t \right) = \frac{1}{c(P, x) \Gamma(m/2 + 1)} (\log(c(P, x)/t))^{m/2}.$$

(ii) Let $\alpha \in P \cap \mathbb{Z}^m$. Then

$$\begin{aligned} \lim_{N \rightarrow \infty} (N/2\pi)^{d(\alpha)/2} D_{N\alpha} \left((N/2\pi)^{d(\alpha)/2} t \right) \\ = \frac{1}{c(P, \alpha) \Gamma(d(\alpha)/2 + 1)} (\log(c(P, \alpha)/t))^{d(\alpha)/2}, \end{aligned}$$

for $0 < t \leq c(P, \alpha)$, where $d(\alpha) = \text{codim } \mu_P^{-1}(\alpha)$.

A sample graph of the scaling limit distribution function for $P = 7\Sigma$ is given in Figure 2.




Figure 2. Scaling limit distribution for $|\varphi_{N\alpha}^P|^2$ with $m = 2, \alpha = (2, 3), P = 7\Sigma$

As in Theorem 1.4, the limit of the rescaled distributions has a universal form, i.e. it does not depend on the geometry of the manifold M_P , and is given by a logarithmic power of the form $(\log c/t)^{d/2}$ with some constant d . The logarithmic power appears because the \mathcal{L}^2 -normalized monomials are close to Gaussian around the peak set $\mu_P^{-1}(x)$ on a vector space of dimension $m/2$ (or dimension $d(\alpha)$ for the boundary lattice cases). More precisely, Theorem 1.2 (and Propositions 3.5 and 3.8) shows that the

function b_x^P has a positive definite Hessian at the peak point. We then observe that the distribution of a Gaussian function on \mathbb{R}^d ,

$$g(u) := \frac{e^{-\langle Au, u \rangle / 2}}{\sqrt{\det A}}, \quad u \in \mathbb{R}^d,$$

(where A is a real positive $(d \times d)$ -matrix) is given by the logarithmic power law

$$\nu_A(u \in \mathbb{R}^d; g(u) > t) = \frac{1}{c\Gamma(d/2 + 1)} \left(\log \frac{c}{t}\right)^{d/2}$$

$$c = \frac{1}{\sqrt{\det A}}, \quad 0 < t \leq c,$$

relative to the normalized Lebesgue measure

$$\nu_A := \frac{\det A}{(2\pi)^{d/2}} du.$$

Thus, the rescaled distribution of an \mathcal{L}^2 -normalized monomial at the ‘center’ of its localized bump has a universal Gaussian form.

To analyze the ‘tails’ of the eigenfunctions, we next use an exponential rescaling of the distribution function so that the global distribution law has a non-zero limit as $N \rightarrow \infty$. As may be expected, it is no longer universal but depends on the geometry of (M_P, ω_P) .

THEOREM 1.5. — *Let $\alpha_N \in NP \cap \mathbb{Z}^m$ satisfy the condition (6) with a point $x \in P^\circ$. Then*

$$\lim_{N \rightarrow \infty} D_{\alpha_N}(e^{-Nt}) = \int_{\{\rho \in \mathbb{R}^m; b_x^P(\rho) < t\}} \det A(\rho) d\rho,$$

where $A(\rho)$ is the Hessian matrix of $\log \sum_{\beta \in P} |c_\beta|^2 e^{\langle \beta, \rho \rangle}$.

A more general scaling limit law is given in Theorem 4.3. In the above theorem, the assumption that $x \in P^\circ$ is not necessary. In fact, we will give similar result for $\alpha_N = N\alpha$ with any lattice point $\alpha \in P$ (Theorem 4.3) in Section 4.

Our strategy for proving Theorems 1.3 and 1.5 on the distribution functions of monomials is based on their pointwise asymptotics (Theorem 1.2, and also Propositions 3.5 and 3.8 in Section 3). Pointwise asymptotics of monomials are more or less equivalent to asymptotics of their L^{2k} norms.

Since the latter are of independent interest, we state these asymptotics explicitly in:

THEOREM 1.6. —

(i) Let $\alpha_N \in NP \cap \mathbb{Z}^m$ satisfy the condition (6) for a point $x \in P^\circ$. Let $\|\varphi_\gamma^P\|_{2k}$ denote the \mathcal{L}^{2k} -norm of the \mathcal{L}^2 -normalized monomial φ_γ^P with the weight $\gamma \in NP$. Then we have

$$\|\varphi_{\alpha_N}\|_{2k}^{2k} = \frac{c(P, x)^{k-1}}{k^{m/2}} \left(\frac{N}{2\pi}\right)^{(k-1)m/2} (1 + O_k(N^{-1})),$$

where $O_k(N^{-1})$ depends on k .

(ii) Let $\alpha \in P \cap \mathbb{Z}^m$ with $d(\alpha) = \text{codim } \mu_P^{-1}(\alpha)$. Then we have

$$\|\widehat{\varphi}_{N\alpha}^P\|_{2k}^{2k} = \frac{c(P, \alpha)^{k-1}}{k^{d(\alpha)/2}} \left(\frac{N}{2\pi}\right)^{(k-1)d(\alpha)/2} (1 + O_k(N^{-1})).$$

We close the introduction with some general remarks and references. As mentioned above, our results pertain to phase space distribution of eigenfunctions (Husimi distributions) rather than to their configuration space distribution. To our knowledge, the only prior example in dimension > 1 for which the limit distribution of eigenfunctions has been determined is the case of certain (so-called) Hecke eigenfunctions of discrete quantum cat maps, due to Kurlberg-Rudnick [KR]. They work in a simpler discrete model rather than the holomorphic model. The main result of Kurlberg-Rudnick [KR] is that the distribution functions of (un-scaled) eigenfunctions tend to the semi-circle law. Their method was to relate the eigenfunctions to exponential sums studied by Katz [Ka] and to apply the value distribution of exponential sums. Since value distribution depends on the representation, it is not clear that the same semi-circle law would hold for Hecke eigenfunctions in the holomorphic (Bargmann-Fock) representation, and this appears to be a challenging and interesting problem. Part of the motivation for this paper was to set a baseline for eigenfunction distribution problems by studying a class of explicitly solvable examples.

It would also be interesting to study the limit distribution of real eigenfunctions in the Schrödinger representation, i.e. on the configuration space rather than the phase space. To our knowledge, the physics results mainly pertain to these configuration space results. The cases most studied

and speculated about are those of chaotic or disordered systems. When eigenfunctions are delocalized, their spatial distribution is conjectured to be Gaussian (see e.g. [B, FE, H, HR,M, SS, PA]). In the opposite regime where the eigenfunctions of disordered systems are exponentially localized, the expected distribution in the low amplitude (tail) region is given by a power of a logarithm [MF], precisely the one we obtained in the high amplitude (center) region. The reason is that the distribution law is universal and Gaussian in the tail region for exponentially localized eigenfunctions of disordered systems [MF], while it is universal and Gaussian in the center for our problem (Theorem 1.2).

The only studies we have located which are related to distribution laws of integrable eigenfunctions are those of one of the authors with J. A. Toth (cf. [TZ]) and that of Berry-Hannay-Ozorio de Almeida [BHO], which describes the asymptotic expansions of \mathcal{L}^{2p} norms (moment intensities) of real oscillatory integrals of several stable types. Such oscillatory integrals define quasimodes for a quantum integrable system, and in generic cases one can express eigenfunctions as oscillatory integrals of various kinds [TZ]. In general, many possible kinds of oscillatory integrals could arise, including ones associated to singular Lagrangean tori. Hence, one can only expect complete results in special cases. To take the simplest example, our methods could be adapted to find the scaling limit distributions of squares of the standard spherical harmonics Y_m^N on S^2 (or S^m). These eigenfunctions are also joint eigenfunctions of commuting operators which generate a quantum torus action. To our knowledge, the distribution laws of even such simple eigenfunctions are unknown at this time.

Acknowledgments: This paper was written during a stay of the second author at Johns Hopkins University on a JSPS fellowship. He would like to express his special thanks to the faculty in the Department of Mathematics of Johns Hopkins University.

2. Background on toric varieties and moment polytopes.

We summarize here some basic facts and terminology on toric varieties from [STZ]. Recall that a *toric variety* is a complex algebraic variety M containing the complex torus

$$(\mathbb{C}^*)^m := (\mathbb{C} \setminus \{0\}) \times \cdots \times (\mathbb{C} \setminus \{0\})$$

as a Zariski-dense open set such that the group action of $(\mathbb{C}^*)^m$ on itself extends to M . We consider here smooth projective toric varieties; they can be given the structure of a symplectic manifold such that the restriction of the action to the underlying real torus

$$\mathbf{T}^m = \{(\zeta_1, \dots, \zeta_m) \in (\mathbb{C}^*)^m : |\zeta_j| = 1, 1 \leq j \leq m\}$$

is a Hamiltonian action (see §2.2). These toric varieties can be constructed from Delzant polytopes either by symplectic reduction (see [Gu]) or by gluing affine toric varieties described by the normal fan of the polytope (see [Fu]). However, for our analysis, it is more convenient to define the toric variety M_P associated to the Delzant polytope P through a *monomial embedding* as follows (see [GKZ Chapter 5]). Suppose that P is a Delzant polytope and let

$$P \cap \mathbb{Z}^m = \{\alpha(1), \alpha(2), \dots, \alpha(d+1)\} .$$

We shall write $\sum_{\alpha \in P} = \sum_{\alpha \in P \cap \mathbb{Z}^m}$. (Recall that a *Delzant polytope* is a convex integral polytope in \mathbb{R}^m with the property that each vertex is incident to exactly m edges and the primitive vectors in \mathbb{Z}^m parallel to these edges generate \mathbb{Z}^m .)

To define the monomial embedding, we fix an arbitrary $c = (c_{\alpha(1)}, \dots, c_{\alpha(d+1)}) \in (\mathbb{C}^*)^{d+1}$. Then, we define

$$\Phi_P^c = [c_{\alpha(1)}\chi_{\alpha(1)}, \dots, c_{\alpha(d+1)}\chi_{\alpha(d+1)}] : (\mathbb{C}^*)^m \rightarrow \mathbb{C}\mathbb{P}^d ; \tag{12}$$

i.e.,

$$\Phi_P^c(z) = [c_{\alpha(1)}z^{\alpha(1)}, \dots, c_{\alpha(d+1)}z^{\alpha(d+1)}], \quad z \in (\mathbb{C}^*)^m .$$

The toric variety $M_P^c = M_P$ is defined as the Zariski-closure of the image $\Phi_P^c((\mathbb{C}^*)^m)$ of the monomial embedding Φ_P^c in the complex projective space $\mathbb{C}\mathbb{P}^d$.

Since our polytope P is assumed to be Delzant, Φ_P^c is an embedding and the variety M_P is smooth. The symplectic (or Kähler) form on M_P is given by

$$(13) \quad \omega_P^c = \omega_P := \Phi_P^{c*} \omega_{\text{FS}} ,$$

where $\omega_{\text{FS}} = \frac{i}{2\pi} \partial\bar{\partial} \log \|\zeta\|^2$ denotes the Fubini-Study Kähler form on $\mathbb{C}\mathbb{P}^d$ with homogeneous coordinates $(\zeta_0, \dots, \zeta_d)$. On $(\mathbb{C}^*)^m$, we have

$$(14) \quad \omega_P^c = \frac{\sqrt{-1}}{2\pi} \partial\bar{\partial} \log \sum_{\alpha \in P} |c_\alpha|^2 |z^\alpha|^2.$$

The volume form on M_P is given by $d\text{Vol}_M = \frac{1}{m!} \omega_P^m$.

2.1. The line bundle L_P^c and associated circle bundle X_P^c .

We define the line bundle $L_P^c \rightarrow M_P^c$ by $L_P = L_P^c := \Phi_P^{c*} \mathcal{O}(1)$, where $\mathcal{O}(1)$ denotes the hyperplane section bundle on $\mathbb{C}\mathbb{P}^d$. Recall that the space of holomorphic sections $H^0(\mathbb{C}\mathbb{P}^d, \mathcal{O}(1))$ consists of the linear functions $\lambda : \mathbb{C}^{d+1} \rightarrow \mathbb{C}$, and that the Fubini-study metric on $\mathcal{O}(1)$ is given by

$$|\lambda|_{\text{FS}}([\zeta]) = \frac{|\lambda(\zeta)|}{\|\zeta\|} \quad (\zeta \in \mathbb{C}^{d+1}),$$

which has curvature form $\omega_{\text{FS}} = \frac{i}{2\pi} \partial\bar{\partial} \log \|\zeta\|^2$. We endow L_P^c with the Hermitian metric $h_P^c := \Phi_P^{c*} h_{\text{FS}}$, which has curvature ω_P^c .

Each monomial χ_α with $\alpha \in P \cap \mathbb{Z}^m$ corresponds to a section of $H^0(M_P^c, L_P^c)$ and vice versa. To explicitly define this correspondence, we make the identifications:

$$(15) \quad \chi_{\alpha(j)}^P \equiv c_{\alpha(j)}^{-1} \Phi_P^{c*} \zeta_j \in H^0(M_P^c, L_P^c) \cong \Phi_P^{c*} H^0(\mathbb{C}\mathbb{P}^d, \mathcal{O}(1)), \quad 1 \leq j \leq d+1.$$

More generally,

$$(16) \quad \Phi_P^{c*} H^0(\mathbb{C}\mathbb{P}^d, \mathcal{O}(N)) \cong H^0(M_P, L_P^N) \quad (N \geq 1),$$

and a basis for $H^0(M_P, L_P^N)$ is given by the sections $\{\chi_\gamma^P : \gamma \in NP \cap \mathbb{Z}^m\}$ corresponding to the monomials $\{\chi_\gamma\}$. These sections are given by

$$\chi_\gamma^P = \chi_{\beta_1}^P \otimes \cdots \otimes \chi_{\beta_N}^P,$$

where $\beta_1, \dots, \beta_N \in P \cap \mathbb{Z}^m$ such that $\gamma = \beta_1 + \cdots + \beta_N$ (see[Fu, STZ]).

So far, we have not specified the constants c_α . For studying our phenomena, the choice of constants defining the toric variety M_P is not important. However, when our polytope P is the full simplex $p\Sigma$, we shall

use the special choice $c_\alpha = \binom{p}{\alpha}^{1/2}$, where $\binom{p}{\alpha}$ is the multinomial coefficient (see §2.4).

The associated principal S^1 -bundle $X_P^c = X_P$ of the line bundle $L_P \rightarrow M_P$ is defined by

$$X_P^c := \{(z, v) \in L_P^{-1}; |v|_P = 1\},$$

where $|v|_P$ denotes the norm of v with respect to the Hermitian metric on L_P^{-1} induced by h_P^c .

We identify sections s_N of L^N with equivariant functions \widehat{s}_N on X by the rule

$$(17) \quad \widehat{s}_N(\lambda) = (\lambda^{\otimes N}, s_N(z)), \quad \lambda \in X_z.$$

Clearly, $\widehat{s}_N(e^{i\theta} \cdot x) = e^{iN\theta} \widehat{s}_N(x)$ if $s_N \in H^0(M_P^c, L_P^N)$. It should be noted that for each $s \in H^0(M_P^c, L_P^N)$, we have

$$|\widehat{s}_N(x)| = |s_N(z)|_P$$

where $x \in X_P$ is in the fiber over $z \in M_P^c$, $|s_N(z)|_P$ denotes the norm with respect to the Hermitian metric on L_P^N induced by the metric h_P^c .

In particular, for $\alpha \in P \cap \mathbb{Z}^m$, the ‘monomial’ $\chi_\alpha^P \in H^0(M_P^c, L_P^c)$ given by (15) lifts to an equivariant function $\widehat{\chi}_\alpha^P$ on the circle bundle $X_P^c \rightarrow M_P^c$, and we write

$$(18) \quad \widehat{m}_{\alpha(j)}^P := c_{\alpha(j)} \widehat{\chi}_{\alpha(j)}^P = \zeta_j \circ \iota_P$$

where $\iota_P : X_P^c \rightarrow S^{2d+1}$ is the lift of the embedding $M_P^c \hookrightarrow \mathbb{C}P^d$ ($d = \#P - 1$), that is, ι_P is the restriction to X_P of the natural inclusion $L_P^{-1} \hookrightarrow \mathcal{O}(-1)$. (Of course, \widehat{m}_α^P depends on c , which we omit to simplify notation.) We also consider the monomials

$$m_\alpha^P := c_\alpha \chi_\alpha^P$$

so that \widehat{m}_α^P is the equivariant lift of m_α to X_P^c . In terms of local coordinates (z, θ) on $\pi^{-1}((\mathbb{C}^*)^m) \subset X_P^c$, we have

$$(19) \quad \widehat{m}_\alpha^P(z, \theta) = \frac{e^{i\theta} c_\alpha z^\alpha}{\left(\sum_{\beta \in P} |c_\beta z^\beta|^2\right)^{1/2}}.$$

Since its absolute value is independent of θ , we can write

$$(20) \quad |m_\alpha^P(z)|_P = |\widehat{m}_\alpha^P(z)| = \frac{|c_\alpha z^\alpha|}{\left(\sum_{\beta \in P} |c_\beta z^\beta|^2\right)^{1/2}}.$$

We give $H^0(M_P, L_P^N)$ the inner product

$$(21) \quad \langle s_1, \bar{s}_2 \rangle = \int_M \langle s_1(z), \overline{s_2(z)} \rangle_{h_N} d\text{Vol}_M(z), \quad s_1, s_2 \in H^0(M_P, L_P^N),$$

and the \mathcal{L}^2 norm $\|s\| = \langle s, \bar{s} \rangle$. We note that the sections

$$\{\chi_\alpha^P \in H^0(M_P, L_P^N) : \alpha = (\alpha_1, \dots, \alpha_m) \in NP\}$$

are orthogonal but not normalized. We normalize them to obtain an orthonormal basis for $H^0(M_P, L_P^N)$ consisting of the sections

$$(22) \quad \varphi_\alpha^P := \frac{\chi_\alpha^P}{\|\chi_\alpha^P\|}.$$

Their equivariant lifts $\widehat{\varphi}_\alpha^P$ form an orthonormal set of monomials on $X_P^\mathbb{C}$.

We include below a table of notation to help the reader keep track of the various monomials:

	monomials on \mathbb{C}^m	sections of L_P^N	monomials on X_P
$\alpha \in NP$	$\chi_\alpha(z) = z^\alpha$	χ_α^P	$\widehat{\chi}_\alpha^P$
$\alpha \in P$		$m_\alpha^P = c_\alpha \chi_\alpha^P \quad (N = 1)$	\widehat{m}_α^P
$\alpha \in NP$		$\varphi_\alpha^P = \chi_\alpha^P / \ \chi_\alpha^P\ $	$\widehat{\varphi}_\alpha^P$

2.2. Moment maps and torus actions.

The group $(\mathbb{C}^*)^m$ acts on $M_P^\mathbb{C}$ and the subgroup \mathbf{T}^m acts in a Hamiltonian fashion. Let us recall the formula for its moment map $\mu_P^\mathbb{C} : M_P^\mathbb{C} \rightarrow \mathbb{R}^m$: on the open orbit $(\mathbb{C}^*)^m$, we have

$$(23) \quad \mu_P(z) = \mu_P^\mathbb{C}(z) = \frac{1}{\sum_{\alpha \in P} |c_\alpha|^2 |z^\alpha|^2} \sum_{\alpha \in P} |c_\alpha|^2 |z^\alpha|^2 \alpha = \sum_{\alpha \in P} |\widehat{m}_\alpha^P(z)|^2 \alpha.$$

For any c , the image of M_P^c under μ_P^c equals P . The moment map μ_P is invariant under the \mathbf{T}^m -action on M_P . By the identification $(\mathbb{C}^*)^m \cong \mathbf{T}^m \times \mathbb{R}^m$, the moment map μ_P defines a map from \mathbb{R}^m to P , which is a diffeomorphism between \mathbb{R}^m and the interior P° of P .

The action of the real torus \mathbf{T}^m lifts from M_P^c to X_P^c and combines with the S^1 action to define a \mathbf{T}^{m+1} action on X_P^c . Recall that under the monomial embedding

$$\Phi_P^c : (\mathbb{C}^*)^m \hookrightarrow M_P^c \hookrightarrow \mathbb{C}\mathbb{P}^d, \quad z \mapsto [c_{\alpha(1)}z^{\alpha(1)}, \dots, c_{\alpha(d+1)}z^{\alpha(d+1)}],$$

the \mathbf{T}^m action on $M_P^c \subset \mathbb{C}\mathbb{P}^d$ is given by

$$(24) \quad e^{i\varphi} \cdot [\zeta_1, \dots, \zeta_{d+1}] = [e^{i\langle \alpha(1), \varphi \rangle} \zeta_1, \dots, e^{i\langle \alpha(d+1), \varphi \rangle} \zeta_{d+1}].$$

The action (24) lifts to an action on L_P^{-1} :

$$(25) \quad e^{i\varphi} \cdot \zeta = (e^{i\langle \alpha(1), \varphi \rangle} \zeta_1, \dots, e^{i\langle \alpha(d+1), \varphi \rangle} \zeta_{d+1}).$$

Since the circle bundle $X_P^c \subset S^{2d+1}$ is invariant under this action, (25) also gives a lift of the action (24) to X_P^c .

We also have the standard circle action on X_P^c :

$$(26) \quad e^{i\theta} \cdot \zeta = e^{i\theta} \zeta,$$

which commutes with the \mathbf{T}^m -action (25). Combining (25) and (26), we then obtain a \mathbf{T}^{m+1} -action on X_P^c :

$$(27) \quad (e^{i\theta}, e^{i\varphi_1}, \dots, e^{i\varphi_m}) \bullet \zeta = e^{i\theta} (e^{i\varphi} \cdot \zeta).$$

2.3. Fourier analysis.

In this section, we shall explain an aspect of Fourier analysis on toric varieties which describe the complete integrability of the system.

The Hardy space $\mathcal{H}^2(X_P^c)$ is the Hilbert space spanned by the equivariant lifts of sections to functions on X_P^c with the inner product

$$\langle f, \bar{g} \rangle = \int_X f \bar{g} dV, \quad dV = \alpha_P \wedge (d\alpha_P)^{n-1},$$

where α_P is the contact 1-form defined by the Hermitian connection on L_P^{-1} such that $d\alpha_P = \pi^*\omega_P$. Under the identification $\mathcal{H} \cong \mathcal{H}^2(X_P^c)$, the inner product is the same as the inner product on $\mathcal{H} = \bigoplus_N H^0(M_P, L_P^N)$ given by (21). Alternately, $\mathcal{H}^2(X_P^c)$ consists of the functions $F \in \mathcal{L}^2(X_P^c)$ satisfying $\bar{\partial}_b F = 0$ (see e.g., [SZ1,Ze]).

Under the S^1 action, the Hardy space then has the orthogonal decomposition

$$(28) \quad \mathcal{H}^2(X_P^c) = \bigoplus_{N=0}^{\infty} \mathcal{H}_N^2(X_P^c),$$

where $\mathcal{H}_N^2(X_P^c)$ consists of elements $\hat{s} \in \mathcal{H}^2(X_P^c)$ such that $\hat{s}(e^{i\theta} \cdot x) = e^{iN\theta} \hat{s}(x)$. We recall that the (equivariant) ‘Szegő projectors’ Π_N are the orthogonal projection onto $H^0(M_P^c, L_P^N) \cong \mathcal{H}_N^2(X_P^c)$. If $\{S_j^N\}$ denotes an orthonormal basis of $H^0(M_P^c, L_P^N)$, and \widehat{S}_j^N denote their lifts to X , then the projector Π_N is given by the kernel function

$$(29) \quad \Pi_N(x, y) = \sum_{j=1}^{k_N} \widehat{S}_j^N(x) \overline{\widehat{S}_j^N(y)} : \mathcal{L}^2(X_P^c) \rightarrow \mathcal{H}_N^2(X_P^c).$$

We now describe how one can combine the eigenvalue problems given by (3) for varying N into a homogeneous scalar eigenvalue problem on X . To do this, we define for each $N \in \mathbb{N}$, the ‘homogenization’ $\widehat{NP} \subset \mathbb{Z}^{m+1}$ of the lattice point in the polytope NP to be the set of all lattice point $\widehat{\alpha}^N$ of the form

$$\widehat{\alpha}^N = \widehat{\alpha} := (Np - |\alpha|, \alpha_1, \dots, \alpha_m), \quad \alpha = (\alpha_1, \dots, \alpha_m) \in NP \cap \mathbb{Z}^m,$$

where $p \geq \max_{\beta \in P \cap \mathbb{Z}^m} |\beta|$. We also define the cone $\Lambda_P = \bigcup_{N=1}^{\infty} \widehat{NP}$. It is well known that rays $\mathbb{N}\widehat{\alpha}$ in this cone define a semiclassical limit.

In this section, we use the more precise notation $\widehat{\varphi}_{\widehat{\alpha}}^P(x)$ for the \mathcal{L}^2 -normalized monomial $\widehat{\varphi}_{\alpha}^P(x)$ (since N is not specified in the latter), for $\widehat{\alpha} \in \Lambda_P$.

The torus action on X_P^c can be quantized to define an action of the torus as unitary operators on $\mathcal{H}^2(X_P^c)$. Specifically, we let $\hat{I}_1, \dots, \hat{I}_m$ denote the differential operators on X_P^c generated by the \mathbf{T}^m action:

$$(30) \quad (\hat{I}_j \hat{S})(\zeta) = \frac{1}{i} \frac{\partial}{\partial \varphi_j} \hat{S}(e^{i\varphi} \cdot \zeta)|_{\varphi=0}, \quad \hat{S} \in \mathcal{C}^{\infty}(X_P^c).$$

We recall the following observation from [STZ]:

PROPOSITION 2.1. — For $1 \leq j \leq m$,

- (i) $\hat{I}_j : \mathcal{H}_N^2(X_P^c) \rightarrow \mathcal{H}_N^2(X_P^c)$;
- (ii) The lifted monomials $\hat{\varphi}_{\hat{\alpha}}^P \in \mathcal{H}_N^2(X_P^c)$ satisfy $\hat{I}_j \hat{\varphi}_{\hat{\alpha}}^P = \alpha_j \hat{\varphi}_{\hat{\alpha}}^P$ ($\hat{\alpha} \in \Lambda_P$).

Furthermore, we note that

$$(31) \quad \frac{\partial}{\partial \theta} : \mathcal{H}_N^2(X_P^c) \rightarrow \mathcal{H}_N^2(X_P^c), \quad \frac{1}{i} \frac{\partial}{\partial \theta} \hat{s}_N = N \hat{s}_N \quad \text{for } \hat{s}_N \in \mathcal{H}_N^2(X_P^c).$$

Thus, the monomials $\hat{\varphi}_{\hat{\alpha}}$ are the solutions of the joint eigenvalue problem

$$(32) \quad \hat{I}_j \hat{\varphi}_{\hat{\alpha}} = \hat{\alpha}_j \hat{\varphi}_{\hat{\alpha}}, \quad \hat{\alpha} \in \mathbb{R}^{m+1}, \quad \bar{\partial}_b \hat{\varphi}_{\hat{\alpha}} = 0, \quad j = 0, \dots, m$$

with the commuting operators:

$$(33) \quad \hat{I}_0 = \frac{p}{i} \frac{\partial}{\partial \theta} - \sum_{j=1}^m \hat{I}_j, \quad \hat{I}_1, \dots, \hat{I}_m.$$

The joint eigenvalues are the lattice points $\hat{\alpha} \in \Lambda_P$.

2.4. Monomials on projective space.

In the case of $\mathbb{C}P^m$, the \mathcal{L}^q -norms of the monomials $\hat{\varphi}_{N\alpha}^{p\Sigma}$ can be evaluated explicitly in an elementary way. We give the details in this section.

When the polytope P is the unit simplex Σ , we have $M_\Sigma = \mathbb{C}P^m$; furthermore L_Σ is the hyperplane bundle $\mathcal{O}(1)$. We can identify $L_\Sigma^{-1} = \mathcal{O}_{\mathbb{C}P^m}(-1)$ with \mathbb{C}^{m+1} with the origin blown up, and the circle bundle $X_\Sigma \subset L_\Sigma^{-1}$ is identified with the unit sphere $S^{2m+1} \subset \mathbb{C}^{m+1}$. The equivariant lifts to X_Σ of sections of $\mathcal{O}(N) = L_\Sigma^N$ consist of homogeneous polynomials

$$F(\zeta_0, \dots, \zeta_m) = \sum_{|\lambda|=N} C_\lambda \zeta^\lambda \quad (\zeta^\lambda = \zeta_0^{\lambda_0} \dots \zeta_m^{\lambda_m})$$

in $m + 1$ variables. The induced Fubini-Study metric on $\mathcal{O}(N)$ is given by

$$|F(\zeta)|_\Sigma = |F(\zeta)| / \|\zeta\|^N, \quad \text{for } F \in H^0(\mathbb{C}P^m, \mathcal{O}(N)).$$

(Here the constants c_α are taken to be 1.) Identifying F with the polynomial $f(z) = F(1, z_1, \dots, z_m)$, the norm can be written

$$|f(z)|_\Sigma = |f(z)| / (1 + \|z\|^2)^{N/2} \quad (z \in \mathbb{C}^m),$$

and the inner product is given by:

$$(34) \quad \langle f, \bar{g} \rangle = \frac{1}{m!} \int_{\mathbb{C}^m} \frac{\langle f(z), \overline{g(z)} \rangle}{(1 + \|z\|^2)^p} \omega_{\text{FS}}^m(z), \quad f, g \in H^0(\mathbb{C}\mathbb{P}^m, \mathcal{O}(p)).$$

We are interested here in the case $P = p\Sigma$. Then $M_{p\Sigma} = \mathbb{C}\mathbb{P}^m$, and the line bundle $L_{p\Sigma}$ is identified with the p -th tensor power $\mathcal{O}(p)$ of the hyperplane section bundle. The circle bundle $X_{p\Sigma}$ is the lens space $X_{p\Sigma} = S^{2m+1}/\mathbb{Z}_p$. By lifting equivariant functions from the lens space $X_{p\Sigma} = S^{2m+1}/\mathbb{Z}_p$ to the sphere S^{2m+1} , we see that $\mathcal{H}_N^2(X_{p\Sigma}) \cong \mathcal{H}_{Np}^2(S^{2m+1})$. Hence, we shall replace the N -th Hardy space $\mathcal{H}_N^2(X_{p\Sigma})$ by $\mathcal{H}_{Np}^2(S^{2m+1})$ below. Then the equivariant lift $\widehat{\chi}_\alpha^{p\Sigma} : S^{2m+1} \rightarrow \mathbb{C}$ of $\chi_\alpha^{p\Sigma} \in H^0(\mathbb{C}\mathbb{P}^m, \mathcal{O}(p))$ is given by the homogenization:

$$\widehat{\chi}_\alpha^{p\Sigma}(x) = x^{\widehat{\alpha}}, \quad \widehat{\alpha} = (p - |\alpha|, \alpha_1, \dots, \alpha_m).$$

In this case, we shall use the special choice of the coefficients of the monomial embedding:

$$c_\alpha^* = \binom{p}{\alpha}^{\frac{1}{2}}, \quad \binom{p}{\alpha} := \frac{p!}{(p - |\alpha|)! \alpha_1! \cdots \alpha_m!},$$

so that by (14), we have

$$\omega_{p\Sigma}^{c^*} = \frac{\sqrt{-1}}{2\pi} \partial\bar{\partial} \log \left(\sum_{|\alpha| \leq p} \binom{p}{\alpha} |z^\alpha|^2 \right) = \frac{\sqrt{-1}}{2\pi} \partial\bar{\partial} \log(1 + \|z\|^2)^p = p\omega_{\text{FS}}.$$

Furthermore,

$$(35) \quad \begin{aligned} \mu_{p\Sigma}(z) &:= \mu_{p\Sigma}^{c^*}(z) = \frac{1}{\sum_{|\alpha| \leq p} \binom{p}{\alpha} |z^\alpha|^2} \sum_{|\alpha| \leq p} \binom{p}{\alpha} |z^\alpha|^2 \alpha \\ &= \frac{p}{1 + \sum |z_j|^2} (|z_1|^2, \dots, |z_m|^2), \end{aligned}$$

where the last equality follows by differentiating the identity $(1 + \sum x_j)^p = \sum_{|\alpha| \leq p} \binom{p}{\alpha} x^\alpha$. Note that this choice gives us the scaling formula

$$\mu_{p\Sigma} = p\mu_\Sigma.$$

As before, for each m -dimensional multi-index β with $|\beta| \leq p$, we define an $(m + 1)$ -dimensional multi-index $\widehat{\beta}$ by

$$\widehat{\beta} = (\widehat{\beta}_0, \widehat{\beta}_1, \dots, \widehat{\beta}_m), \quad \widehat{\beta}_0 = p - |\beta|, \quad \widehat{\beta}_j = \beta_j \quad (j = 1, \dots, m).$$

Recall that $|\widehat{\varphi}_\beta^{p\Sigma}(x)| = |\varphi_\beta^{p\Sigma}(z)|_{p\Sigma}$ with $x \in S^{2m+1}$, $\pi(x) = z \in \mathbb{C}\mathbb{P}^m$.

The \mathcal{L}^q -norms of the monomial $\widehat{\varphi}_{N\widehat{\alpha}}^{p\Sigma}$ can be evaluated explicitly as follows.

PROPOSITION 2.2. — For $\alpha \in p\Sigma \cap \mathbb{Z}^m$, we have the precise formula:

$$\|\widehat{\varphi}_{N\widehat{\alpha}}^{p\Sigma}\|_q^q = \left[\frac{(Np + m)!}{(N\widehat{\alpha})!} \right]^{q/2} \frac{\prod_{j=0}^m \Gamma(Nq\widehat{\alpha}_j/2 + 1)}{p^{m(q/2-1)} \Gamma(Npq/2 + m + 1)}.$$

Proof. — We write $\widehat{\chi}_{N\widehat{\alpha}} = \widehat{\chi}_{N\widehat{\alpha}}^{p\Sigma}$, which we consider as a function in $\mathcal{H}_{Np}^2(S^{2m+1})$. For $q \geq 1$, we set

$$\mathcal{I}_q(N) = \int_{\mathbb{C}^{m+1}} e^{-|x|^2} |\widehat{\chi}_{N\widehat{\alpha}}(x)|^q d\ell(x),$$

where $d\ell(x)$ denotes Lebesgue measure on \mathbb{C}^{m+1} . We shall compute the integral $\mathcal{I}_q(N)$ in two ways. First we use polar coordinates on \mathbb{C}^{m+1} . The measure $d\ell$ is expressed as

$$d\ell(x) = r^{2m+1} dr d\sigma, \quad x = r\sigma, \quad r > 0, \quad \sigma \in S^{2m+1}.$$

Then we have

$$\begin{aligned} \mathcal{I}_q(N) &= \int_0^\infty e^{-r^2} r^{Npq+2m+1} dr \int_{S^{2m+1}} |\widehat{\chi}_{N\widehat{\alpha}}(\sigma)|^q d\sigma \\ &= \frac{1}{2} \Gamma(Npq/2 + m + 1) \int_{S^{2m+1}} |\widehat{\chi}_{N\widehat{\alpha}}(\sigma)|^q d\sigma. \end{aligned}$$

To relate the volume form $d\sigma$ on S^{2m+1} to that on $X_{p\Sigma}$, we recall that $\omega_{p\Sigma} = p\omega_{\text{FS}}$ and therefore the volume form on $X_{p\Sigma}$ is p^m times the Fubini-Study volume. Recalling our convention that $\text{Vol}(X_\Sigma) = \text{Vol}(\mathbb{C}\mathbb{P}^m) = \frac{1}{m!}$, we then have $d\text{Vol}_{X_{p\Sigma}} = \frac{p^m}{2\pi^{m+1}} d\sigma$, and hence

$$\mathcal{I}_q(N) = \frac{\pi^{m+1}}{p^m} \Gamma(Npq/2 + m + 1) \|\widehat{\chi}_{N\hat{\alpha}}\|_q^q.$$

On the other hand, if we use polar coordinates for each component x_j of $x \in \mathbb{C}^{m+1}$, we have

$$\begin{aligned} \mathcal{I}_q(N) &= \prod_{j=0}^m \int_{\mathbb{C}} e^{-|x|^2} |x|^{N\alpha_j q} d\ell(x) \\ &= (2\pi)^{m+1} \prod_{j=0}^m \int_0^\infty e^{-r^2} r^{N\alpha_j q+1} dr = \pi^{m+1} \prod_{j=0}^m \Gamma(N\alpha_j q/2 + 1). \end{aligned}$$

Hence we obtain

$$(36) \quad \|\widehat{\chi}_{N\hat{\alpha}}\|_q^q = p^m \frac{\prod_{j=0}^m \Gamma(Nq\alpha_j/2 + 1)}{\Gamma(Npq/2 + m + 1)}$$

and therefore

$$(37) \quad \|\widehat{\chi}_{N\hat{\alpha}}\|^2 = \frac{p^m (N\hat{\alpha})!}{(Np + m)!}.$$

Since $\widehat{\varphi}_{N\hat{\alpha}}^{p\Sigma} = \widehat{\chi}_{N\hat{\alpha}} / \|\widehat{\chi}_{N\hat{\alpha}}\|$, the identity follows from (36)–(37). □

The following proposition is a direct consequence of Stirling’s formula and Proposition 2.2.

PROPOSITION 2.3. — *Let $\alpha \in p\Sigma \cap \mathbb{Z}^m$, and let $J = \{j \in \mathbb{Z} : 0 \leq j \leq m, \alpha_j \neq 0\}$, where $\alpha_0 = p - |\alpha|$. Set $d(\alpha) = 2m - \#J$. Then we have*

$$\|\widehat{\varphi}_{N\hat{\alpha}}^{p\Sigma}\|_q^{2q} \sim \left(\frac{N}{2\pi}\right)^{(q/2-1)d(\alpha)} \frac{p^{q/2-1} (2\pi)^{r(q-2)}}{(q/2)^{d(\alpha)} \left(\prod_{j \in J} \alpha_j\right)^{q/2-1}};$$

$$\|\widehat{\varphi}_{N\hat{\alpha}}^{p\Sigma}\|_\infty^2 = p^{-m} \frac{(Np + m)!}{(N\hat{\alpha})!} \left[\frac{\prod_{j \in J} \alpha_j^{\alpha_j}}{p^p} \right]^N.$$

In particular, again by Stirling’s formula, we have

$$\|\widehat{\varphi}_{N\alpha}^{p\Sigma}\|_{\infty}^2 \sim (2\pi)^r \left(\frac{p}{\prod_{j \in J} \alpha_j}\right)^{1/2} \left(\frac{N}{2\pi}\right)^{d(\alpha)/2}.$$

In the next section, we will obtain similar formulas for monomials on general toric varieties.

3. Pointwise asymptotics on general toric varieties.

We now consider the case of general toric varieties. Our first purpose is to find the pointwise asymptotics of the monomials and to prove Theorem 1.2. We then asymptotically determine their \mathcal{L}^{2k} -norms.

3.1. Pointwise asymptotics: interior points.

First of all, we shall consider a sequence α_N of lattice points in NP . We assume that α_N is an approximate multiple of a point $x \in P^\circ$ as in Definition 1.1; i.e.,

$$(38) \quad \alpha_N/N = x + O(N^{-1}).$$

Hence, the point α_N/N is in the interior P° of the polytope P for sufficiently large N . Thus the analysis is performed on the open orbit $(\mathbb{C}^*)^m$, and hence the coordinate

$$z = e^{\rho/2+i\varphi} \in (\mathbb{C}^*)^m, \quad \rho, \varphi \in \mathbb{R}^m$$

will be useful. The moment map μ_P is also invariant under the Hamiltonian \mathbf{T}^m -action, and it is well-known ([Fu]) that it induces a diffeomorphism:

$$(39) \quad \bar{\mu}_P : \mathbb{R}^m = (\mathbb{C}^*)^m/\mathbf{T}^m \rightarrow P^\circ, \quad \bar{\mu}_P(\rho) := \mu_P(e^{\rho/2}).$$

In these coordinates, the function b_x^P defined by (7) can be written simply as

$$(40) \quad b_x^P(\rho) = f(x, \rho) - f(x, \rho_x^P), \quad \rho_x^P = \bar{\mu}_P^{-1}(x), \quad f(x, \rho) := \log k(\rho) - \langle \rho, x \rangle,$$

where the function $k(\rho)$ is the ‘polytope character’

$$(41) \quad k(\rho) = \sum_{\beta \in P \cap \mathbb{Z}^m} |c_\beta|^2 e^{\langle \rho, \beta \rangle}.$$

The vector $\tau_x^P(z)$ in (8) is given by

$$(42) \quad \tau_x^P(z) = \rho - \rho_x^P, \quad z = e^{\rho/2 + i\theta}.$$

We also define the symmetric real $m \times m$ matrix

$$(43) \quad A(P, x) := \sum_{\beta \in P} \left| \widehat{m}_\beta^P(e^{\rho_x^P/2}) \right|^2 \beta \otimes \beta - x \otimes x.$$

LEMMA 3.1. — *The real symmetric matrix*

$$A(\rho) := \sum_{\beta \in P} |\widehat{m}_\beta^P(e^{\rho/2})|^2 \beta \otimes \beta - \mu_P(e^{\rho/2}) \otimes \mu_P(e^{\rho/2})$$

is positive definite, for all $z \in (\mathbb{C}^*)^m$.

Proof. — We must show that $(\lambda \otimes \lambda, A(\rho)) > 0$ for $\lambda \in (\mathbb{R}^m)' \setminus \{0\}$. Consider the vectors $u, v \in \mathbb{R}^{d+1}$ given by $u_\alpha = |\widehat{m}_\alpha^P(e^{\rho/2})|$, $v_\alpha = |\widehat{m}_\alpha^P(e^{\rho/2})| \lambda(\alpha)$. Since $\|u\|^2 = \sum_{\alpha \in P} |\widehat{m}_\alpha^P(e^{\rho/2})|^2 = 1$ and $\mu_P(e^{\rho/2}) = \sum_{\alpha \in P} |\widehat{m}_\alpha^P(e^{\rho/2})|^2 \alpha$, we have

$$(44) \quad \begin{aligned} (\lambda \otimes \lambda, A(\rho)) &= \sum_{\alpha \in P} |\widehat{m}_\alpha^P(e^{\rho/2})|^2 \lambda(\alpha)^2 - \left(\sum_{\alpha \in P} |\widehat{m}_\alpha^P(e^{\rho/2})|^2 \lambda(\alpha) \right)^2 \\ &= \|v\|^2 - \langle u, v \rangle^2 = \|u\|^2 \|v\|^2 - \langle u, v \rangle^2 \geq 0. \end{aligned}$$

Since $u_\alpha = |\widehat{m}_\alpha^P(e^{\rho/2})| \neq 0$ for all $\alpha \in P \cap \mathbb{Z}^m$ and $v_\alpha/u_\alpha = \lambda(\alpha)$ is not constant on $P \cap \mathbb{Z}^m$, the Cauchy-Schwartz inequality in (44) is strict. \square

In particular, the matrix

$$A(P, x) = A(\rho_x^P)$$

is positive definite.

For a point $x \in P^\circ$, we now define the constant

$$(45) \quad c(P, x) := \frac{1}{\sqrt{\det A(P, x)}}.$$

LEMMA 3.2. — *In the coordinates $z = e^{\rho/2+i\theta}$ on $(\mathbb{C}^*)^m$, the volume form on M_P is given by*

$$\omega_P^m/m! = \frac{1}{(2\pi)^m} \det A(\rho) d\rho d\theta.$$

Proof. — We note that

$$(46) \quad A(\rho) = \sum_{\beta \in P} k_\beta(\rho) \beta \otimes \beta - \left(\sum_{\beta \in P} k_\beta(\rho) \beta \right) \otimes \left(\sum_{\beta \in P} k_\beta(\rho) \beta \right) = \text{Hess}_\rho \log k(\rho),$$

where $k_\beta(\rho) = |\widehat{m}_\beta^P(e^{\rho/2})|^2$ and $k(\rho) = \sum_{\beta \in P} |c_\beta|^2 e^{\langle \rho, \beta \rangle}$. The conclusion follows from (46), recalling that

$$(47) \quad \omega_P = \Phi_P^* \omega_{\text{FS}} = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log \sum_{\beta} |c_\beta|^2 |\chi_\beta(z)|^2 = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log k(\rho). \quad \square$$

It should be noted that $b_x^P(\rho)$ grows as $|\rho| \rightarrow \infty$, as stated in the following simple lemma.

LEMMA 3.3. — *Let $K \subset P^\circ$ be a compact set. Then there exists positive constants $R > 0, c > 0$ such that $f(x, \rho) \geq c|\rho|$ for $(x, \rho) \in K \times \mathbb{R}^m, |\rho| \geq R$*

Proof. — For any $(x, \rho) \in P^\circ \times \mathbb{R}^m$, we define

$$M(x, \rho) = \max_{\beta \in P \cap \mathbb{Z}^m} \langle \rho, \beta - x \rangle.$$

If $x \in P^\circ$, then the polytope $P - x$ contains the origin in its interior. Thus, clearly we have $M(x, \rho) > 0$ for any $(x, \rho) \in P^\circ \times (\mathbb{R}^m \setminus \{0\})$. Next, we note that the function $(x, \rho) \mapsto M(x, \rho)$ is continuous. To see this, let (x_n, ρ_n) be a sequence such that $(x_n, \rho_n) \rightarrow (x, \rho) \in P^\circ \times \mathbb{R}^m$. Then, for any $\beta \in P \cap \mathbb{Z}^m$,

$$|\langle \rho_n, \beta - x_n \rangle - \langle \rho, \beta - x \rangle| \leq C|\rho_n - \rho| + |\rho||x_n - x|.$$

By using this inequality, we can show that

$$|M(x_n, \rho_n) - M(x, \rho)| \leq C|\rho_n - \rho| + |\rho||x_n - x|.$$

Now, for a compact set $K \subset P^o$, we set

$$M(K) = \min_{(x, \rho) \in K \times \mathbb{R}^m, |\rho|=1} M(x, \rho) > 0.$$

We set $c_0 = \min_{\beta \in P \cap \mathbb{Z}^m} |c_\beta|^2$. Since $P \cap \mathbb{Z}^m$ is a finite set, there exists $\beta = \beta(x, \rho)$ such that $M(x, \rho) = \langle \rho, \beta(x, \rho) - x \rangle$ for (x, ρ) with $x \in P^o$ and $|\rho| = 1$. Thus, for $x \in K \subset P^o$ and $\rho \neq 0$, we have

$$e^{f(x, \rho)} = \sum_{\beta} |c_\beta|^2 e^{\langle \rho, \beta - x \rangle} \geq c_0 e^{|\rho| \langle \frac{\rho}{|\rho|}, \beta(x, \frac{\rho}{|\rho|}) - x \rangle} = c_0 e^{|\rho| M(x, \frac{\rho}{|\rho|})} \geq c_0 e^{|\rho| M(K)}$$

for $x \in K$ and $\rho \neq 0$, which completes that proof. □

Completion of the proof of Theorem 1.2: Recalling that $z = e^{\rho/2+i\theta}$, we write $|\varphi_{\alpha_N}^P(\rho)|_P$ instead of $|\varphi_{\alpha_N}^P(z)|_P$. By the definition of the Hermitian metric on L_P^N , we have

$$|\varphi_{\alpha_N}^P(\rho)|_P^2 = \frac{|\chi_{\alpha_N}^P(\rho)|_P^2}{\|\chi_{\alpha_N}^P\|^2} = \frac{1}{\|\chi_{\alpha_N}^P\|^2} \frac{e^{\langle \rho, \alpha_N \rangle}}{|\Phi_P^c(e^{\rho/2})|^{2N}},$$

where, in the right hand side, $|\Phi_P^c(e^{\rho/2})|$ denotes the usual norm in \mathbb{C}^{d+1} . By the definition (12) of the monomial embedding Φ_P^c , we have

$$|\Phi_P^c(e^{\rho/2})|^2 = k(\rho), \quad \rho \in \mathbb{R}^m.$$

Hence, we have

$$(48) \quad |\varphi_{\alpha_N}^P(\rho)|_P^2 = \frac{e^{-Nf(\alpha_N/N, \rho)}}{\|\chi_{\alpha_N}^P\|^2} = \frac{e^{-Nf(x, \rho)} R_N(x, \rho)}{\|\chi_{\alpha_N}^P\|^2}, \quad R_N(x, \rho) = e^{N\langle \rho, \alpha_N/N - x \rangle},$$

where the function $f(x, \rho)$ for $x \in P^o$ is defined in (40). We note that, since $\alpha_N/N - x = O(N^{-1})$, we have

$$(49) \quad |\partial_\rho^L R_N(x, \rho)| \leq C_L R_N(x, \rho)$$

for every multi-index L . By Lemma 3.2, the \mathcal{L}^2 -norm of the un-normalized monomial $\chi_{\alpha_N}^P$ is given by

$$\|\chi_{\alpha_N}^P\|^2 = \int_{\mathbb{R}^m} e^{-Nf(\alpha_N/N, \rho)} \det A(\rho) d\rho.$$

Here it should be noted that $\det A(\rho)$ is a positive integrable function on \mathbb{R}^m . By Lemma 3.3, we can choose $R > 0, c > 0$ such that $|\rho_x^P| < R$ and $f(\alpha_N/N, \rho) \geq c|\rho|$ for any $|\rho| \geq R$ and N . Thus, by choosing a cut-off function $g(\rho)$ suitably, we may write

$$\|\chi_{\alpha_N}^P\|^2 = e^{-Nf(x, \rho_x^P)} \int e^{-Nb_x^P(\rho)} R_N(x, \rho) g(\rho) \det A(\rho) d\rho + O(e^{-cN}).$$

Recall that $b_x^P(\rho) = 0$ if and only if $\rho = \rho_x^P$, and that $\rho = \rho_x^P$ is the unique critical point of b_x^P . The Hessian of b_x^P at $\rho = \rho_x^P$ is the positive definite symmetric matrix $A(P, x)$. Thus, by the Morse lemma, there exists a change of coordinates κ from a neighborhood of the origin to a neighborhood of ρ_x^P such that $\kappa(0) = \rho_x^P$ and that

$$b_x^P \circ \kappa(\xi) = \langle A(P, x)\xi, \xi \rangle / 2, \quad |\det D\kappa(0)| = 1.$$

By choosing the cut-off function g suitably, we get

$$\int e^{-Nb_x^P(\rho)} g(\rho) R_N(x, \rho) \det A(\rho) d\rho = \int e^{-N\langle A(P, x)\xi, \xi \rangle / 2} G_N(x, \xi) d\xi,$$

where $G_N(x, \xi)$ is a compactly supported function in ξ such that

$$G_N(x, 0) = R_N(x, \rho_x^P) \det A(P, x),$$

and the function $R_N(x, \rho)$ is defined in (48). By (49), the derivatives of $G_N(x, \xi)$ with respect to ξ are all bounded uniformly in N . Therefore, by using the Plancherel formula and a formula for the Fourier transform of the Gaussian functions, we obtain

$$\begin{aligned} \int e^{-Nb_x^P(\rho)} g(\rho) R_N(x, \rho) \det A(\rho) d\rho \\ = \left(\frac{N}{2\pi}\right)^{-m/2} R_N(x, \rho_x^P) \sqrt{\det A(P, x)} (1 + O(N^{-1})), \end{aligned}$$

which, combined with (48), completes the proof. □

Remark. — Since $\chi_{\alpha_N}^P$ is a monomial, the asymptotics of $\|\chi_{\alpha_N}^P\|$ is essentially the same calculation as the asymptotics of the L^{2k} norm of another monomial. Thus, determining the pointwise asymptotics of monomials is equivalent to determining the asymptotics of their L^{2k} norms.

3.2. Pointwise asymptotics: boundary lattice points.

Next, we consider the ray $\mathbb{N}\alpha$ for a lattice point $\alpha \in P \cap \mathbb{Z}^m$, which is allowed to lie in the boundary ∂P . In such a case, we need to work with other coordinates than the usual coordinates on the open orbit $(\mathbb{C}^*)^m$ ([SZ2]), since the open orbit $(\mathbb{C}^*)^m$ does not cover the set $\mu_P^{-1}(\partial P)$.

In the following, we mean that the faces are disjoint, and the facet is a face of codimension one. Thus we call the closed face (or facet) \bar{F} the closure of F in the minimal affine subspace containing F . To describe the coordinates, let v_0 be a vertex of P . Since our polytope P is Delzant, we can choose lattice points $\alpha^1, \dots, \alpha^m$ in P such that each α^j is in an edge incident to the vertex v_0 , and the vectors $v^j := \alpha^j - v_0$ form a basis of \mathbb{Z}^m . We choose (open) facets $F_j, j = 1, \dots, m$ incident at v_0 so that $\alpha^j \notin F_j$.

LEMMA 3.4. — *Let $\alpha \in P \cap \mathbb{Z}^m$, and $z \in M_P$. Then, $\chi_\alpha^P(z) = 0$ if and only if*

$$(50) \quad \mu_P(z) \in \bigcup \{ \bar{F} ; F \text{ is a facet } \alpha \notin \bar{F} \}.$$

Proof. — If $z \in (\mathbb{C}^*)^m$ then automatically we have $\chi_\alpha^P(z) \neq 0$. Thus, we may assume that $z \in \mu_P^{-1}(\partial P)$. First, assume that $\mu_P(z) \in \bar{F}$ for some closed facet \bar{F} which does not contain α . The last formula in (23) for the moment map is globally defined, for the function $|\widehat{m}_\beta^P|^2$ is globally defined. Since $\mu_P(z) \in \bar{F}$, the coefficients in $\mu_P(z)$ of the lattice points $\beta \notin \bar{F}$ must vanish. Thus, we have $\chi_\alpha^P(z) = 0$. Conversely, assume that $\mu_P(z) \in \partial P$ is not in the set described in (50). In our convention, the faces are disjoint and the boundary ∂P is the disjoint union of faces. Thus, that $\mu_P(z) \in \partial P$ is not in the set described in (50) is equivalent to say that there exists an open face E such that $\alpha \in \bar{E}$ and $\mu_P(z) \in E$. Let v_1, \dots, v_l be the set of vertex of \bar{E} . Since v_j and α are lattice points, there exists a positive integer n_0 such that $n_0\alpha = \sum_{j=1}^l n_j v_j$ with n_j integer such that $\sum n_j = n_0$. Thus we have

$$(\chi_\alpha^P)^{\otimes n_0}(z) = (\chi_{v_1}^P)^{\otimes n_1}(z) \otimes \dots \otimes (\chi_{v_l}^P)^{\otimes n_l}(z).$$

Since $\mu_P(z)$ is in the interior E of the face (polytope) \bar{E} , each $\chi_{v_j}^P(z)$ can not vanish, and hence $\chi_\alpha^P(z) \neq 0$. □

To apply Lemma 3.4, we set

$$U_{v_0} := \{z \in M_P ; \chi_{v_0}^P(z) \neq 0\},$$

which covers M_P as v_0 varies over all vertices. We define

$$(51) \quad \eta : (\mathbb{C}^*)^m \rightarrow (\mathbb{C}^*)^m, \quad \eta(z) = \eta_j(z) := (z^{v^1}, \dots, z^{v^m}).$$

The map η is a diffeomorphism and the inverse is given by

$$z : (\mathbb{C}^*)^m \rightarrow (\mathbb{C}^*)^m, \quad z(\eta) = (\eta^{\Gamma e^1}, \dots, \eta^{\Gamma e^m}),$$

where e^j is the standard basis for \mathbb{R}^m (or for \mathbb{C}^m over \mathbb{C}), and Γ is an $m \times m$ -matrix with $\det \Gamma = \pm 1$ and integer coefficients defined by

$$\Gamma v^j = e^j, \quad v^j = \alpha^j - v_0.$$

By definition, we have the obvious formula:

$$\chi_{\alpha^j}^P(z) = \eta_j(z) \chi_{v_0}^P(z), \quad z \in (\mathbb{C}^*)^m.$$

By Lemma 3.4, $\eta_j(z) \rightarrow 0$ if $z \in U_{v_0}$, $z \rightarrow \mu_P^{-1}(\bar{F}_j)$. Since $\alpha^j \notin \bar{F}_j$, we have

$$(\chi_{\alpha^j}^P)^{-1}(0) \cap U_{v_0} = \mu_P^{-1}(\bar{F}_j).$$

The set $U_{v_0} \setminus (\mathbb{C}^*)^m$ is the union of the sets $\mu_P^{-1}(\bar{F}_j)$, and hence the map η extends a homeomorphism:

$$\eta : U_{v_0} \rightarrow \mathbb{C}^m, \quad \eta(z_0) = 0, \quad z_0 = \text{the fixed point corresponding to } v_0.$$

By this homeomorphism, the set $\mu_P^{-1}(\bar{F}_j)$ corresponds to the set $\{\eta \in \mathbb{C}^m; \eta_j = 0\}$. This coordinate $\eta = (\eta_1, \dots, \eta_m)$ is useful to explain toric subvarieties corresponding to faces. Namely, let \bar{F} be a closed face with $\dim F = m - r$ which contains v_0 . Since $v_0 \in \bar{F}$, we can choose F_{i_1}, \dots, F_{i_r} such that $\bar{F} = \bar{F}_{i_1} \cap \dots \cap \bar{F}_{i_r}$. Then the subvariety $\mu_P^{-1}(\bar{F})$ corresponding \bar{F} is expressed, in the coordinate neighborhood U_{v_0} , by

$$\mu_P^{-1}(\bar{F}) \cap U_{v_0} = \{\eta \in \mathbb{C}^m; \eta_{i_j} = 0, \quad j = 1, \dots, r\}.$$

Now, we fix a lattice point α in a (relatively open) face F of dimension $\dim F = m - r$ such that $v_0 \in \bar{F}$. Without loss of generality, we may assume that $\bar{F} = \bar{F}_1 \cap \dots \cap \bar{F}_r$.

To state a result for the lattice point α in the boundary corresponding to Theorem 1.2, we need to find a function corresponding to the function b_α^P .

Since our coordinate η is based on the lattice points $\alpha^j - v_0$, it is reasonable to introduce a new polytope defined by the affine linear transformation

$$\tilde{\Gamma} : \mathbb{R}^m \ni u \rightarrow \Gamma u - \Gamma v_0 \in \mathbb{R}^m,$$

which maps \mathbb{Z}^m bijectively onto itself. We set $Q := \tilde{\Gamma}(P)$. Then Q is contained in the positive orthant $\{x \in \mathbb{R}^m; x_j \geq 0\}$, and we have $\tilde{\Gamma}(\bar{F}_j) = \{x \in Q; x_j = 0\}$. The face of Q corresponding to F is then given by

$$Q_F := \tilde{\Gamma}(\bar{F}) = \{x \in Q; x_j = 0, \quad j = 1, \dots, r\}.$$

We denote a point in $U_{v_0} \cong \mathbb{C}^m$ as $\eta = (\xi, \zeta) \in \mathbb{C}^m = \mathbb{C}^r \times \mathbb{C}^{m-r}$. In this expression, $\zeta = (0, \zeta)$ is a local coordinate of the submanifold $\mu_P^{-1}(\bar{F})$. The modulus square of the monomials $|\chi_{N\alpha}^P|^2$ with $\alpha \in P \cap \mathbb{Z}^m$ is, in this coordinate, given by

$$(52) \quad |\chi_{N\alpha}^P(z)|^2 = \frac{|\eta^{\tilde{\Gamma}(\alpha)}|^{2N}}{K(\eta)^N},$$

$$K(\eta) = \sum_{\gamma \in Q \cap \mathbb{Z}^m} a_\gamma |\eta^\gamma|^2, \quad a_\gamma = |c_{\tilde{\Gamma}^{-1}(\gamma)}|^2.$$

We then introduce the ‘moment map’ corresponding to the face F by:

$$(53) \quad \mu_F : \mathbb{R}^{m-r} \rightarrow Q_F, \quad \mu_F(\rho) = \sum_{(0, \nu) \in Q_F} \frac{a_\nu e^{\langle \rho, \nu \rangle}}{k_F(\rho)} (0, \nu),$$

where $a_\nu = |c_{\tilde{\Gamma}^{-1}(0, \nu)}|^2$, and the function $k_F(\rho)$ is given by

$$(54) \quad k_F(\rho) := \sum_{(0, \nu) \in Q_F} a_\nu e^{\langle \rho, \nu \rangle}.$$

As mentioned above, the submanifold $\mu_P^{-1}(\bar{F})$ has the coordinate $\zeta \mapsto (0, \zeta) \in \mathbb{C}^r \times \mathbb{C}^{m-r}$, and the torus \mathbf{T}^{m-r} acts on it. Thus, it is natural to use the coordinate

$$\zeta = e^{\rho/2 + i\theta}, \quad \rho, \theta \in \mathbb{R}^{m-r}.$$

Then, we write $\eta = (\xi, \zeta) = (\xi, \rho)$ for $\zeta = e^{\rho/2}$. Since we have assumed $\alpha \in F$, we may write

$$\tilde{\Gamma}(\alpha) = (0, \tilde{\alpha}) \in Q_F.$$

We also define

$$s_\alpha(\xi, \rho) := \log K(\xi, \rho) - \langle \rho, \tilde{\alpha} \rangle = \log(k_F(\rho) + \ell_F(\xi, \rho)) - \langle \rho, \tilde{\alpha} \rangle, \tag{55}$$

$$\ell_F(\xi, \rho) = \sum_{(\mu, \nu) \in Q, \mu \neq 0} a_{(\mu, \nu)} |\xi^\mu|^2 e^{\langle \rho, \nu \rangle}.$$

PROPOSITION 3.5. — *In the coordinate $\eta = (\xi, \zeta)$ as above, we write $|\varphi_{N\alpha}^P(\xi, \rho)|_P$ for the modulus square of the monomial $|\varphi_{N\alpha}^P|_P^2$. We also write $\tilde{\Gamma}(\alpha) = (0, \tilde{\alpha})$, which is in the interior of the polytope Q_F . Then we have*

$$|\varphi_{N\alpha}^P(\xi, \rho)|_P^2 = (2\pi)^r \left(\frac{N}{2\pi} \right)^{(m+r)/2} \frac{e^{-N\Psi_\alpha(\xi, \rho)}}{\sqrt{\det A(F, \alpha)}} (1 + O(N^{-1})),$$

where the function $\Psi_\alpha(\xi, \rho)$ is given by

$$\Psi_\alpha(\xi, \rho) = s_\alpha(\xi, \rho) - s_\alpha(0, \rho_\alpha^F) \quad \rho_\alpha^F = \mu_F^{-1}(\tilde{\alpha}) \in \mathbb{R}^{m-r}. \tag{56}$$

and $(m - r) \times (m - r)$ positive definite matrix $A(F, \alpha)$ is given by

$$A(F, \alpha) = \sum_{(0, \nu) \in Q_F} \frac{a_\nu e^{\langle \rho_\alpha^F, \nu \rangle}}{k_F(\rho_\alpha^F)} \nu \otimes \nu - \tilde{\alpha} \otimes \tilde{\alpha}. \tag{57}$$

We shall prove Proposition 3.5 in the rest of this subsection. First of all, we need the following simple lemma:

LEMMA 3.6. — *In the coordinate $(\xi, \zeta = e^{\rho/2+i\theta})$ on $\mathbb{C}^r \times (\mathbb{C}^*)^{m-r} \subset U_{v_0}$, the volume form $\omega_P^m/m!$ is given by*

$$\frac{\omega_P^m}{m!} = \frac{1}{\pi^r (2\pi)^{m-r}} L(\xi, \rho) dm(\xi) d\rho d\theta, \tag{58}$$

where $dm(\xi)$ denotes the Lebesgue measure on \mathbb{C}^r , and the function $L(\xi, \rho)$ is given by the determinant of the following $m \times m$ matrix:

$$L(\xi, \rho) = \det \begin{pmatrix} \frac{\partial^2 \log K}{\partial \xi \partial \xi} & \frac{\partial^2 \log K}{\partial \xi \partial \rho} \\ \frac{\partial^2 \log K}{\partial \rho \partial \xi} & \frac{\partial^2 \log K}{\partial \rho \partial \rho} \end{pmatrix}, \tag{59}$$

where $K(\xi, \rho)$ is the function defined in (52).

Proof. — For the function $f(\xi, \zeta)$ on $\mathbb{C}^r \times (\mathbb{C}^*)^{m-r}$ independent of the variable θ in $\zeta = e^{\rho/2+i\theta}$, then the derivatives $(\partial f)/(\partial \zeta_j)$, $(\partial f)/(\partial \bar{\zeta}_j)$ is given, respectively, by

$$\frac{\partial f}{\partial \zeta_j} = \frac{1}{\zeta_j} \frac{\partial f}{\partial \rho_j}, \quad \frac{\partial f}{\partial \bar{\zeta}_j} = \frac{1}{\bar{\zeta}_j} \frac{\partial f}{\partial \rho_j}.$$

We write $\eta = (\xi, \zeta)$. Then, by this relation, we have

$$\det \left(\frac{\partial^2 \log K}{\partial \eta_j \partial \bar{\eta}_k} \right) = \left(\prod_{j=1}^{m-r} |\zeta_j|^2 \right)^{-1} L(\xi, \rho),$$

where $L(\xi, \rho)$ is given by (59). But, we have

$$dm(\zeta) = \frac{1}{2^{m-r}} \left(\prod_{j=1}^{m-r} |\zeta_j|^2 \right) d\rho d\theta,$$

where $dm(\zeta)$ denotes the Lebesgue measure on \mathbb{C}^{m-r} . Combining this with (47), we obtain the assertion. □

The following lemma can be shown by the same argument as in the proof of Lemma 3.1, and we shall omit the proof.

LEMMA 3.7. — *The $(m - r) \times (m - r)$ matrix defined by*

$$A_F(\rho) := \partial \mu_F(\rho) = \sum_{(0, \nu) \in Q_F} \frac{a_\nu e^{\langle \rho, \nu \rangle}}{k_F(\rho)} \nu \otimes \nu - \mu_F(\rho) \otimes \mu_F(\rho)$$

is positive definite for every $\rho \in \mathbb{R}^{m-r}$.

Note that, the map $\mu_F : \mathbb{R}^{m-r} \rightarrow Q_F^o$ is a diffeomorphism, and the lattice point $\tilde{\alpha}$ is in Q_F^o . Thus, the vector $\rho_\alpha^F = \mu_F^{-1}(\tilde{\alpha})$ is well-defined. Hence, the $(m - r) \times (m - r)$ matrix

$$A(F, \alpha) = A_F(\rho_\alpha^F)$$

is positive definite.

the polytope $Q = \tilde{\Gamma}(P)$ contains the standard basis in $\mathbb{R}^m = \mathbb{R}^r \times \mathbb{R}^{m-r}$. Thus the lattice points $(e_j^r, 0)$ with the standard basis e_j^r in \mathbb{R}^r is in the polytope Q , and hence the functions f_j are all positive. This combined with Lemma 3.7 shows that the Hessian $Hs_\alpha(0, \rho_\alpha)$ is positive definite. By the same argument as in the proof of Lemma 3.3, the function $k_F(\rho) + \ell_F(\xi, \rho)$ tends to ∞ as $|\xi| + |\rho| \rightarrow \infty$. Therefore, by the standard Laplace method as in the proof of Theorem 1.2, we have

$$(63) \quad \|\chi_{N\alpha}^P\|^2 = \frac{1}{\pi^r} \left(\frac{N}{2\pi}\right)^{-(m+r)/2} \frac{e^{-Ns_\alpha(0, \rho_\alpha^F)}}{\sqrt{\det Hs_\alpha(0, \rho_\alpha^F)}} L(0, \rho_\alpha^F)(1 + O(N^{-1})).$$

A direct computation will show that

$$(64) \quad \det Hs_\alpha(0, \rho_\alpha^F) = \frac{\det A(F, \alpha)}{k_F(\rho_\alpha^F)^{2r}} \left(\prod_{j=1}^r 2f_j(\rho_\alpha^F)\right)^2,$$

$$L(0, \rho) = \frac{\det A_F(\rho)}{k_F(\rho_\alpha^F)^r} \left(\prod_{j=1}^r f_j(\rho)\right),$$

and hence, the asymptotics (63) can be written in the form:

$$\|\chi_{N\alpha}^P\|^2 = \frac{1}{(2\pi)^r} \left(\frac{N}{2\pi}\right)^{-(m+r)/2} \sqrt{\det A(F, \alpha)} e^{-Ns_\alpha(0, \rho_\alpha^F)}(1 + O(N^{-1})).$$

Dividing $|\chi_{N\alpha}|_P^2$ by the above, we conclude the assertion. □

When our fixed lattice point α is a vertex, say $\alpha = v_0$ in the description of the coordinate η , the matrix $A(F, \alpha)$ is not defined suitably. However, clearly the similar asymptotics can be deduced by the same method.

PROPOSITION 3.8. — *Suppose that α is a vertex of the polytope P . Then we have*

$$|\varphi_{N\alpha}^P(\eta)|_P^2 = \left(\frac{N}{|c_\alpha|^2}\right)^m e^{-N(\log K(\eta) - \log |c_\alpha|^2)}(1 + O(N^{-1})),$$

where the function $K(\eta)$ on \mathbb{C}^m is given by (52). We also have $|c_\alpha|^2 = |\chi_\alpha^P(z_\alpha)|^{-2}$ where z_α is the fixed point for the Hamiltonian \mathbf{T}^m -action such that $\mu_P(z_\alpha) = \alpha$.

Proof. — In the description of the coordinate η , we put $\alpha = v_0$. Then, clearly we have $K(0) = |c_\alpha|^2 = |\chi_\alpha^P(z_\alpha)|^{-2}$, where the fixed point z_α

corresponds to the origin $\eta = 0$ and the function $K(\eta)$ is defined in (52). In this case, we just use the coordinate η itself without change of variable. In this coordinate, the monomial $|\chi_{N\alpha}^P|_P^2$ is given by

$$|\chi_{N\alpha}^P(\eta)|_P^2 = e^{-N \log K(\eta)}.$$

The volume measure $\omega_P^m/m!$ is of the form:

$$\frac{\omega_P^m}{m!} = \frac{1}{\pi^m} \det L(\eta) dm(\eta), \quad L(\eta) = \left(\frac{\partial^2 \log K(\eta)}{\partial \eta_j \partial \bar{\eta}_k} \right)$$

with the Lebesgue measure $dm(\eta)$ on \mathbb{C}^m . It is straightforward to see that the critical point of the function $\log K(\eta)$ is the origin, and the determinant of the Hessian at the origin is given by

$$H(\log K)(0) = \begin{pmatrix} \frac{2L(0)}{K(0)} & 0 \\ 0 & \frac{2L(0)}{K(0)} \end{pmatrix}.$$

By using these facts with the Laplace method, we obtain the assertion. \square

3.3. Moments and \mathcal{L}^{2k} norms.

As an application of the pointwise estimates, one can prove that eigenfunctions ‘localize on tori’. We also determine \mathcal{L}^{2k} norms of the monomials. Before proceeding with the discussion on asymptotics of \mathcal{L}^{2k} norms, we need the following simple lemma, which asserts that, in view of the pointwise asymptotics (Theorem 1.2), monomials decay exponentially as $N \rightarrow \infty$, away from the invariant torus $\mu_P^{-1}(x)$.

LEMMA 3.9. — *Let x be a point in the interior of the polytope P , and let $\alpha_N \in NP \cap \mathbb{Z}^m$ be an approximate multiple of x . Then, there exists positive constants $c > 0$ and $R > 0$ such that we have*

$$b_x^P(\rho) - \langle \rho - \rho_x^P, x_N \rangle \geq c|\rho|, \quad x_N = \alpha_N/N - x, \quad |\rho| \geq R.$$

Proof. — By using Lemma 3.3 and the definition of the function b_x^P , there exists constants $c_0 > 0$, $R_0 > 0$ such that

$$b_x^P(\rho) - \langle \rho - \rho_x, x_N \rangle \geq c_0|\rho| - \langle \rho, x_N \rangle + A_x,$$

where we set $A_x = -f(x, \rho_x^P) + \langle \rho_x^P, x_N \rangle$. We note that $x_N \rightarrow 0$ as $N \rightarrow \infty$. Thus, we only need to choose $c > 0$ and $R > 0$ so that $c_0 - |x_N| > c$ for sufficiently large N and $R \geq A_x / (c_0 - |x_N| - c)$. \square

The following is easily shown by using Theorem 1.2, Propositions 3.5, Lemma 3.9 and 3.8, and the argument is the same as in their proofs, and hence we shall omit the proof (see also the proof of Theorem 3.12).

PROPOSITION 3.10. —

(i) Let $\alpha_N \in NP \cap \mathbb{Z}$ be an approximate multiple of a point $x \in P^\circ$. Then, the measure $|\varphi_{\alpha_N}^P|_P^2 d\text{Vol}_{M_P}$ weak*-converges to the normalized Haar measure on the m -dimensional torus $\mu_P^{-1}(x)$, i.e.,

$$\int_{M_P} \sigma |\varphi_{\alpha_N}^P|_P^2 d\text{Vol}_{M_P} \rightarrow \int_{\mu_P^{-1}(x)} \sigma d\theta$$

for $\sigma \in \mathcal{C}(M_P)$.

(ii) Let α be a lattice point in P with $\dim \mu_P^{-1}(\alpha) = m - r$. Then, we have

$$w^* \lim_{N \rightarrow \infty} |\varphi_{N\alpha}^P|_P^2 = d\theta_{\mu_P^{-1}(\alpha)},$$

where $d\theta_{\mu_P^{-1}(\alpha)}$ denotes the normalized Haar measure on the $(m - r)$ -dimensional torus $\mu_P^{-1}(\alpha) \cong \mathbf{T}^{m-r}$.

In the above proposition, if α is a vertex, then the left hand side denotes the Dirac measure at the fixed point $z_\alpha \in M_P$ of the Hamiltonian \mathbf{T}^m -action corresponding to α .

We note that $|\widehat{\varphi}_{N\alpha}^P|$ is invariant under the \mathbf{T}^m action. We denote the Hilbert space of \mathbf{T}^m invariant functions by $\mathcal{L}_{\text{inv}}^2(M_P)$. We can restate the conclusion as follows: if $\sigma \in C_{\text{inv}}^\infty(M_P)$, then we can regard it as a function on the polytope P . We can also regard $|\widehat{\varphi}_{N\alpha}^P|^2$ as a function, say $|\widetilde{\varphi}_\alpha^P(I)|^2$, on P , equipped with action variables I . We then have:

COROLLARY 3.11. — For any $\alpha \in P \cap \mathbb{Z}^m$, we have $|\widetilde{\varphi}_{N\alpha}^P|^2 dI \rightarrow \delta_\alpha$; i.e., $\int_P \sigma |\widetilde{\varphi}_\alpha^P|^2 dI \rightarrow \sigma(\alpha)$, for $\sigma \in \mathcal{C}(P)$. For $\alpha_N \in NP \cap \mathbb{Z}^m$ satisfying (38) with a point $x \in P^\circ$, we have $|\widetilde{\varphi}_{\alpha_N}^P|^2 dI \rightarrow \delta_x$.

Next, we determine the asymptotics of the \mathcal{L}^{2k} -norm of the \mathcal{L}^2 -normalized monomials.

THEOREM 3.12. —

(i) Let $\alpha_N \in NP \cap \mathbb{Z}^m$ be an approximate multiple of a point $x \in P^o$. Let $\|\varphi_\gamma^P\|_{2k}$ denote the \mathcal{L}^{2k} -norm of the \mathcal{L}^2 -normalized monomial φ_γ^P with the weight $\gamma \in NP \cap \mathbb{Z}^m$. Then we have

$$(65) \quad \|\varphi_{\alpha_N}^P\|_{2k}^{2k} = \frac{1}{k^{m/2}} \left(\frac{N}{2\pi}\right)^{(k-1)m/2} \frac{1}{(\det A(P, x))^{(k-1)/2}} (1 + O_k(N^{-1})),$$

where the $O_k(N^{-1})$ depends on k .

(ii) Let $\alpha \in P$ be a lattice point with $\dim \mu_P^{-1}(\alpha) = m - r$. Then, for $r \leq m - 1$, we have

$$(66) \quad \|\widehat{\varphi}_{N\alpha}^P\|_{2k}^{2k} = \frac{1}{k^{(m+r)/2}} \left(\frac{N}{2\pi}\right)^{(k-1)(m+r)/2} \left(\frac{(2\pi)^r}{\sqrt{\det A(F, \alpha)}}\right)^{k-1} (1 + O_k(N^{-1})).$$

For a vertex α , we have

$$(67) \quad \|\widehat{\varphi}_{N\alpha}^P\|_{2k}^{2k} = \frac{1}{k^m} \left(\frac{N}{|c_\alpha|^2}\right)^{(k-1)m} (1 + O_k(N^{-1})).$$

Proof. — The proof of (67) is the same as that for (66), so, first of all, we shall give a proof of (66). By Theorem 1.2 and Proposition 3.5, we have

$$\begin{aligned} \|\widehat{\varphi}_{N\alpha}^P\|_{2k}^{2k} &= \frac{(2\pi)^{kr}}{\pi^r} \left(\frac{N}{2\pi}\right)^{k(m+r)/2} \frac{1}{(\det A(F, \alpha))^{k/2}} \\ &\quad \int_{\mathbb{C}^r \times \mathbb{R}^{m-r}} e^{-Nk\Psi_\alpha(\xi, \rho)} L(\xi, \rho) dm(\xi) d\rho (1 + O_k(N^{-1})), \end{aligned}$$

where Ψ_α is given by (56). As in the proof of Proposition 3.5, the critical point of Ψ_α is only the point $(0, \rho_\alpha^F)$ with $\rho_\alpha^F = \mu_F^{-1}(\tilde{\alpha})$, $\tilde{\Gamma}(\alpha) = (0, \tilde{\alpha})$, and which is non-degenerate. We have $\Psi_\alpha(0, \rho_\alpha^F) = 0$. Thus, by the standard Laplace method, we have

$$\begin{aligned} &\|\widehat{\varphi}_{N\alpha}^P\|_{2k}^{2k} \\ &= \frac{(2\pi)^{kr}}{\pi^r} \left(\frac{N}{2\pi}\right)^{(k-1)(m+r)/2} \frac{1}{(\det A(F, \alpha))^{k/2}} \frac{L(0, \rho_\alpha)}{\det H\Psi_\alpha(0, \rho_\alpha)} (1 + O_k(N^{-1})). \end{aligned}$$

Therefore, the assertion follows from (64) as in the proof of Proposition 3.5.

To prove (65), we use Theorem 1.2 and Lemma 3.9 to find

$$\begin{aligned} & \|\varphi_{\alpha_N}^P\|_{2k}^{2k} \\ &= c(P, x)^k \left(\frac{N}{2\pi}\right)^{km/2} \int e^{-Nkb_x^P(\rho)} R_N(\rho) \det A(\rho) f(\rho) d\rho (1 + O(N^{-1})) \end{aligned}$$

where we set $R_N(\rho) = e^{N\langle \rho - \rho_x^P, x_N \rangle}$ and f is a cut-off function around ρ_x^P . Note that the function $R_N(\rho)$ and its derivatives are dominated from above by the function $R_N(\rho)$ itself, since $Nx_N = O(1)$. Thus, as in the proof of Theorem 1.2, we obtain the asymptotics (65) by using the standard Laplace method. \square

By using Proposition 3.5, we can determine the limit of the sup-norm $\|\varphi_{N\alpha}^P\|_{\infty}^2$.

To do this, we prepare the following lemma, which shows that, when $x \in P^\circ$ and γ_N is an approximate multiple of x , the monomials $|\varphi_{\gamma_N}^P|^2$ look like Gaussian on a ball around the invariant torus $\mu_P^{-1}(x)$ of radius $O(N^{-1/2})$.

LEMMA 3.13. — *Let x be a point in the interior of the polytope P , and let α_N be an approximate multiple of x . For any positive number r , we set*

$$B_x(r) = \{z = e^{\rho/2+i\theta} \in (\mathbb{C}^*)^m; |\rho - \rho_x^P| < r\},$$

which is a neighborhood of the invariant torus $\mu_P^{-1}(x)$ corresponding to x . Let c be a positive number. Then we have

$$(68) \quad |\varphi_{\alpha_N}^P(z)|^2 = \left(\frac{N}{2\pi}\right)^{m/2} \frac{e^{-\langle A(P,x)u, u \rangle/2}}{\sqrt{\det A(P,x)}} (1 + O(N^{-1/2})),$$

$$z = e^{(\rho_x^P + u/\sqrt{N})/2+i\theta} \in B_x(c/\sqrt{N}), \quad |u| \leq c.$$

Proof. — Taylor expansion for the function b_x^P around the point ρ_x^P shows

$$(69) \quad b_x^P(\rho_x^P + v) = \int_0^1 (1-t) \langle A(\rho_x^P + tv)v, v \rangle dt = \frac{1}{2} \langle A(P, x)v, v \rangle + O(|v|^3).$$

Now, let $z = e^{(\rho_x^P + u/\sqrt{N})/2 + i\theta} \in B_x(c/\sqrt{N})$ with $|u| \leq c$. Then, we have

$$b_x^P(z) = \frac{1}{2N} \langle A(P, x)u, u \rangle + O(N^{-3/2}).$$

For such a $z \in B_x(c/\sqrt{N})$, we also have

$$\langle \tau_x^P(z), x_N \rangle = O(N^{-3/2}),$$

because $x_N = O(1/N)$. Thus, the Taylor expansion concludes the assertion. □

PROPOSITION 3.14. —

(i) Let $\alpha_N \in NP \cap \mathbb{Z}^m$ satisfy (38) with a point $x \in P^\circ$. Then we have

$$(70) \quad \lim_{N \rightarrow \infty} \left(\frac{N}{2\pi} \right)^{m/2} \|\varphi_{\alpha_N}^P\|_\infty^2 = \frac{1}{\sqrt{\det A(P, x)}}.$$

(ii) Let α be a lattice point with $\dim \mu_P^{-1}(\alpha) = m - r$, $r \leq m - 1$. Let (ξ_N, ρ_N) be the point where $|\varphi_{N\alpha}^P|_P^2$ attains its maximum. Then, we have

$$(71) \quad \lim_{N \rightarrow \infty} \left(\frac{N}{2\pi} \right)^{-(m+r)/2} \|\varphi_{N\alpha}^P\|_\infty^2 = \frac{(2\pi)^r}{\sqrt{\det A(F, \alpha)}}.$$

For a vertex $\alpha \in P$, we have

$$(72) \quad \lim_{N \rightarrow \infty} N^{-m} \|\varphi_{N\alpha}^P\|_\infty^2 = |c_\alpha|^{-2m} = |\chi_\alpha^P(z_\alpha)|_P^{2m},$$

where $z_\alpha \in M_P$ is the unique fixed point for the \mathbf{T}^m such that $\mu_P(z_\alpha) = \alpha$.

Proof. — First of all, we shall show (71). The function $\Psi_\alpha(\xi, \rho)$ defined in (56) attains its minimum at the point $(0, \rho_\alpha^F)$, and it tends to ∞ as $|\xi| + |\rho| \rightarrow \infty$. Thus, the function $(\frac{N}{2\pi})^{-(m+r)/2} |\varphi_{N\alpha}^P|_P^2$ is of order $O(e^{-cN})$ outside a compact neighborhood of $(0, \rho_\alpha^F)$. On a compact neighborhood B of $(0, \rho_\alpha^F)$, we have

$$|\varphi_{N\alpha}^P(0, \rho_\alpha^F)|_P^2 \leq \sup_{(\xi, \rho) \in B} |\varphi_{N\alpha}^P(\xi, \rho)|_P^2 \leq \left(\frac{N}{2\pi} \right)^{(m+r)/2} \frac{(2\pi)^r}{\sqrt{\det A(F, \alpha)}} (1 + O(N^{-1})).$$

Now (71) follows from the above inequality. The same argument with Proposition 3.5 shows (72).

Next, we shall prove (70). By Lemma 3.9, the function $|\varphi_{\alpha_N}^P(\rho)|^2$ attains its maximum on a compact set around the point ρ_x^P . We set $f_N(\rho) := (\frac{N}{2\pi})^{-m/2}|\varphi_{\alpha_N}(\rho)|^2$, and let $\rho_N = \rho_x^P + v_N$ be the point such that $f_N(\rho_N) = (\frac{N}{2\pi})^{-m/2}\|\varphi_{\alpha_N}\|_\infty^2$ with $|v_N| \leq c$ for some constant $c > 0$ independent of N . Then, by Theorem 1.2, we have

$$b_x(\rho_N) - \langle \rho_N - \rho_x^P, x_N \rangle = O(1/N).$$

Since $x_N = O(N^{-1})$, we have $0 \leq b_x^P(\rho_N) = O(N^{-1})$. By the Taylor expansion for the function b_x^P ((69)), we can choose a constant $a > 0$ such that

$$b_x^P(\rho_x^P + v) \geq a|v|^2, \quad |v| \leq c.$$

Thus, we obtain $|v_N| = O(N^{-1/2})$. We set $v_N = u_N/\sqrt{N}$ so that $|u_N| \leq c$. Then, by Lemma 3.13, we have

$$f_N(\rho_x^P) \leq f_N(\rho_x^P + u_N/\sqrt{N}) = \frac{1}{\sqrt{\det A(P, x)}} e^{-\langle A(P, x)u_N, u_N \rangle/2} (1 + O(N^{-1})),$$

and hence

$$e^{\langle A(P, x)u_N, u_N \rangle/2} \leq f_N(\rho_x^P)^{-1} \frac{1}{\sqrt{\det A(P, x)}} (1 + O(N^{-1})) \rightarrow 1$$

as $N \rightarrow \infty$. This shows that $u_N \rightarrow 0$ as $N \rightarrow \infty$. Therefore, we have $f_N(\rho_x^P + u_N/\sqrt{N}) \rightarrow 1/\sqrt{\det A(P, x)}$ as $N \rightarrow \infty$. \square

3.4. Asymptotics on projective space.

The values of the \mathcal{L}^{2k} norm in Proposition 2.2 and in the projective-space case of Theorem 1.6 may seem to be different. However, we can check that these two coincide by noting the following simple lemma.

LEMMA 3.15. — *The determinant $\det A(p\Sigma, \alpha)$ of the matrix $A(p\Sigma, \alpha)$ is given by*

$$\det A(p\Sigma, \alpha) = \frac{(p - |\alpha|)\alpha_1 \cdots \alpha_m}{p}.$$

Proof. — By applying the differential operator $x_j \partial_{x_j}$ twice to the formula $(1 + \sum_{l=1}^m x_l)^p = \sum_{|\beta| \leq p} \binom{p}{\beta} x^\beta$ for a vector $x \in \mathbb{R}^m$, we have

$$p(p-1)x_i x_j (1 + \sum_l x_l)^{p-2} + p x_j (1 + \sum_l x_l)^{p-1} \delta_{ij} = \sum_{|\beta| \leq p} \binom{p}{\beta} x^\beta \beta_i \beta_j,$$

where δ_{ij} is the Kronecker's delta, and we have set $\binom{p}{\beta} = p! / \beta! (p - |\beta|)!$. We put $r = \sum e^{(\rho_\alpha)_j}$. Then, by (35), we have $e^{\rho_\alpha} = \frac{1+r}{p} \alpha$ and $|\widehat{n}_\beta^{p\Sigma}(e^{\rho_\alpha/2})|^2 = e^{\langle \rho_\alpha, \beta \rangle} \binom{p}{\beta} / (1+r)^p$. From this the second equation follows. Substituting $\frac{1+r}{p} \alpha$ for x in the above formula, we obtain

$$A(p\Sigma, \alpha)_{i,j} = \sum_{|\beta| \leq p} \binom{p}{\beta} \frac{e^{\langle \rho, \beta \rangle}}{(1+r)^p} \beta_i \beta_j - \alpha_i \alpha_j = \alpha_j \delta_{ij} - \frac{1}{p} \alpha_i \alpha_j.$$

We set $D_m(\alpha_1, \dots, \alpha_m) = \det A(p, \alpha)$ with $\alpha = (\alpha_1, \dots, \alpha_m)$. Then a simple computation shows that

$$\frac{D_m(\alpha_1, \dots, \alpha_m)}{\alpha_1 \cdots \alpha_m} = \frac{D_{m-1}(\alpha_2, \dots, \alpha_m)}{\alpha_2 \cdots \alpha_m} - \frac{1}{p} \alpha_1.$$

Thus the lemma follows by induction on the dimension m . □

Note that, for the simplex $p\Sigma$, the faces containing the origin is again a simplex in lower dimensional vector space. Therefore, Lemma 3.15 can be applied to compute $\det A(F, \alpha)$.

In the case of projective monomials, the function $b_\alpha^{p\Sigma}$ can be expressed as follows.

PROPOSITION 3.16. — For $z \in (\mathbb{C}^*)^m$,

$$b_\alpha^{p\Sigma}(z) = p \log(1 + |z|^2) - \log |z^\alpha|^2 + \langle \widehat{\alpha}, \log \widehat{\alpha} \rangle - p \log p,$$

where $\log \widehat{\alpha} = (\log \widehat{\alpha}_0, \dots, \log \widehat{\alpha}_m)$.

This is a direct consequence of the formulas $e^{\rho_\alpha} = \frac{1+r}{p} \alpha$, $r = \sum e^{(\rho_\alpha)_j}$ and $(1 + \sum_l x_l)^p = \sum_{|\beta| \leq p} \binom{p}{\beta} x^\beta$ as mentioned before.

The pointwise asymptotics of the \mathcal{L}^2 -normalized projective monomials is given in the following proposition.

PROPOSITION 3.17. —

(1) For $z \in (\mathbb{C}^*)^m$,

$$|\widehat{\varphi}_{N\alpha}^{p\Sigma}(z)|^2 = \left(\frac{N}{2\pi}\right)^{m/2} \frac{p^{1/2} e^{-N b_\alpha^{p\Sigma}(z)}}{\sqrt{(p - |\alpha|)\alpha_1 \cdots \alpha_m}} (1 + O(N^{-1})),$$

uniformly $(\mathbb{C}^*)^m$.

(2) For $z = e^{(\rho_\alpha + u/\sqrt{N})/2 + i\theta}$, we have

$$|\widehat{\varphi}_{N\alpha}^{p\Sigma}(z)|^2 = \left(\frac{N}{2\pi}\right)^{m/2} \frac{p^{1/2} e^{-\langle \Delta(\alpha)u, u \rangle - \frac{1}{p} \langle \alpha, u \rangle^2} / 2}{\sqrt{(p - |\alpha|)\alpha_1 \cdots \alpha_m}} (1 + O(N^{-1/2}))$$

uniformly for $|u| \leq c$, where $\Delta(\alpha)$ is the diagonal matrix with entries $\alpha_1, \dots, \alpha_m$.

The assertion (1) in the above is a restatement of Theorem 1.2 for the projective monomials. The assertion (2) follows from (1) and a Taylor expansion of the function $b_\alpha^{p\Sigma}$.

4. Asymptotics of distribution functions.

In this section, we find asymptotics of rescaled and un-rescaled distribution functions. Fix a lattice point α in P , and let r denote the codimension of the face of P (possibly the open face P°) containing α . In analogy with (45), we define the constant $c(P, \alpha)$ by

$$(73) \quad c(P, \alpha) := \begin{cases} \frac{(2\pi)^r}{\sqrt{\det A(F, \alpha)}} & \text{if } r < m \\ \frac{(2\pi)^m}{|c_\alpha|^{2m}} = (2\pi |\chi_{N\alpha}^P(z_\alpha)|_P^2)^m & \text{if } \alpha \text{ is a vertex.} \end{cases}$$

In the following discussion, we give the details for the case where $\alpha_N = N\alpha$ with a lattice point $\alpha \in P \cap \mathbb{Z}^m$. For general $\alpha_N \in NP \cap \mathbb{Z}^m$ satisfying (38) for a point $x \in P^\circ$, one needs only to put $r = 0$ and replace $N\alpha$ and α by α_N and x .

4.1. Rescaled distribution functions.

We would like to understand the limit distribution of the measures $|\widehat{\varphi}_{N\alpha}^P|_*^2 d\text{Vol}_{M_P}$, namely, the limit of their distribution functions

$$(74) \quad D_{N\alpha}(t) := \text{Vol}_{M_P} \{z; |\widehat{\varphi}_{N\alpha}^P(z)|^2 > t\}.$$

However, by Theorem 1.6, the k -th moments of the measure $|\widehat{\varphi}_{N\alpha}^P|_*^2 d\text{Vol}_{M_P}$ tends to infinity as N tends to infinity. Therefore, we need to re-normalize the monomials.

We write

$$(75) \quad dv_N^r = \left(\frac{N}{2\pi}\right)^{(m+r)/2} d\text{Vol}_{M_P}, \quad f_{N\alpha} = \left(\frac{N}{2\pi}\right)^{-(m+r)/4} \varphi_{N\alpha}^P(z)$$

so that

$$\int_{M_P} |f_{N\alpha}(z)|_P^2 dv_N^r(z) = 1.$$

By Proposition 3.14, we know that $\lim_{N \rightarrow \infty} \|f_{N\alpha}\|_\infty^2$ exists and we have

$$(76) \quad \lim_{N \rightarrow \infty} \|f_{N\alpha}\|_\infty^2 = c(P, \alpha).$$

Furthermore, by Theorem 1.6, we have

$$(77) \quad \|f_{N\alpha}\|_{\mathcal{L}^{2k}(dv_N^r)}^{2k} = \left(\frac{N}{2\pi}\right)^{-(m+r)(k-1)/2} \|\widehat{\varphi}_{N\alpha}^P\|_{2k}^{2k} = \frac{c(P, \alpha)^{k-1}}{k^{(m+r)/2}} (1 + O(N^{-1})).$$

We consider the limit distribution of the sequence of measures

$$(78) \quad \nu_{N,r} := |f_{N\alpha}|_*^2 dv_N^r$$

on the real line. The distribution function $F_{N\alpha}^r(t)$ of the measure $\nu_{N,r}$ is given by

$$(79) \quad F_{N\alpha}^r(t) = \left(\frac{N}{2\pi}\right)^{(m+r)/2} \text{Vol}_{M_P} \{z \in M_P; |\widehat{\varphi}_{N\alpha}^P(z)|^2 > \left(\frac{N}{2\pi}\right)^{(m+r)/2} t\}.$$

The distribution function $F_{N\alpha}^r$ defined above can be expressed, in terms of the distribution function $D_{N\alpha}$ for the measure $|\widehat{\varphi}_{N\alpha}^P|_*^2 d\text{Vol}_{M_P}$, as

$$F_{N\alpha}^r(t) = \left(\frac{N}{2\pi}\right)^{(m+r)/2} D_{N\alpha} \left(\left(\frac{N}{2\pi}\right)^{(m+r)/2} t \right).$$

We should note that the total mass of the measure $\nu_{N,r}$ is $\text{Vol}(M_P)(N/2\pi)^{(m+r)/2}$, and hence it tends to infinity as N goes to infinity. But its k -th moment satisfies, for each positive integer k ,

$$(80) \quad \int x^k d\nu_{N,k}(x) = \frac{c(P, \alpha)^{k-1}}{k^{(m+r)/2}} (1 + O_k(N^{-1})) \rightarrow \frac{c(P, \alpha)^{k-1}}{k^{(m+r)/2}} \quad (N \rightarrow \infty).$$

By Proposition 3.14, the support of the measure $\nu_{N,r}$ is contained in a bounded interval in $[0, +\infty)$ which is independent of N .

Furthermore, by (80), the measure $x d\nu_{N,r}(x)$ is a finite measure on the real line whose support lies in a bounded interval in $[0, +\infty)$ which is independent of N . This implies that the sequence of the finite measures $x d\nu_{N,r}(x)$ on the real line has a weak limit, say μ , and it must satisfy

$$(81) \quad \int x^k d\mu(x) = \frac{c(P, \alpha)^k}{(k + 1)^{(m+r)/2}}, \quad k = 0, 1, 2, \dots$$

Thus, the weak limits of the measures $x d\nu_{N,r}(x)$ are probability measures, and they have supports contained in the interval $[0, c(P, \alpha)]$.

LEMMA 4.1. — *Let c be a positive constant, and let h be a positive integer. Let μ be a probability measure on the real line such that*

$$\int x^k d\mu(x) = \frac{c^k}{(k + 1)^{h/2}}, \quad k \geq 0.$$

Then μ must coincide with the measure $\rho_{c,h}(x) dx$ where

$$(82) \quad \rho_{c,h}(x) = \frac{1}{c\Gamma(h/2)} \chi_{(0,c)}(x) (\log(c/x))^{h/2-1}.$$

Proof. — We shall use the following formula:

$$\frac{1}{w^s} = \frac{1}{\Gamma(s)} \int_0^\infty e^{-wt} t^{s-1} dt$$

to compute the Fourier transform

$$\hat{\mu}(\xi) = \int e^{ix\xi} d\mu(x)$$

of the measure μ satisfying the condition in the lemma. Substituting $s = h/2, w = k$ in the above formula, we get

$$\begin{aligned} \hat{\mu}(\xi) &= \sum_{k \geq 0} \frac{(ic\xi)^k}{k!(k+1)^{h/2}} = \frac{1}{\Gamma(h/2)} \int_0^\infty e^{ic\xi e^{-t}} e^{-t} t^{h/2-1} dt \\ &= \frac{1}{c\Gamma(h/2)} \int_0^c e^{i\xi x} (\log(c/x))^{h/2-1} dx, \end{aligned}$$

and hence $\hat{\mu}(\xi) = \hat{\rho}_{c,h}(\xi)$. This completes the proof. □

As a corollary to Lemma 4.1, we have the following.

COROLLARY 4.2. — *The sequence of the finite measures $\{x d\nu_{N,r}(x)\}$ converges weakly to the probability measure $\rho_{c(P,\alpha),r} dx$, where the density $\rho_{c(P,\alpha),r}$ is given by (82) with $c = c(P, \alpha)$ and $h = m + r$.*

Proof of Theorem 1.4. The rescaled distribution function $F_{N\alpha}^r(t)$ for $t > 0$ of the measure $\nu_N = |f_{N\alpha}|_*^2 d\nu_N^r$ is given by

$$F_{N\alpha}^r(t) = \int \chi_{(t,+\infty)}(x) d\nu_{N,r}(x),$$

where $\chi_{(t,+\infty)}$ is the characteristic function of the interval $(t, +\infty)$. Now set $d\mu_{N,r}(x) = x d\nu_{N,r}(x)$ and write

$$F_{N\alpha}^r(t) = \int x^{-1} \chi_{(t,+\infty)}(x) d\mu_{N,r}(x).$$

By Corollary 4.2, we know that the sequence of finite measures μ_N converges weakly to the probability measure $\rho_{c(P,\alpha),r}(x) dx$. For fixed $t > 0$, by approximating the function $x^{-1} \chi_{(t,+\infty)}(x)$ by a sequence of continuous functions, and by using the Lebesgue convergence theorem, one easily obtains that, for every $0 < t \leq c(P, \alpha)$,

$$\begin{aligned} \lim_{N \rightarrow +\infty} F_{N\alpha}^r(t) &= \int x^{-1} \chi_{(t,+\infty)}(x) \rho_{c(P,\alpha),r}(x) dx \\ &= \frac{1}{c(P, \alpha)\Gamma((m+r)/2)} \int_t^{c(P,\alpha)} x^{-1} (\log(c(P, \alpha)/x))^{(m+r)/2-1} dx \\ &= \frac{1}{c(P, \alpha)\Gamma((m+r)/2)} \int_0^{\log(c(P,\alpha)/t)} s^{(m+r)/2-1} ds \\ &= \frac{1}{c(P, \alpha)\Gamma((m+r)/2 + 1)} (\log(c(P, \alpha)/t))^{(m+r)/2}. \end{aligned} \quad \square$$

4.2. Non-rescaled distribution functions.

In the previous section, we derived the limit of the rescaled distribution functions $F_{N\alpha}^r(t)$. In the definition of the rescaled function $F_{N\alpha}^r(t)$, the parameter t is rescaled by the factor $(N/2\pi)^{(m+r)/2}$ so that it tends to $+\infty$ as $N \rightarrow \infty$. Then the corresponding volume

$$(83) \quad \text{Vol}_{M_P}(z; |\widehat{\varphi}_{N\alpha}^P(z)|^2 > (N/2\pi)^{(m+r)/2}t)$$

is of order $N^{-(m+r)/2}$. This was the reason why we need to multiply the volume (83) by the extra factor $(N/2\pi)^{(m+r)/2}$ in the definition of the rescaled distribution functions. As a result, the limit distribution has a universal form (Theorem 1.4). However, one may ask, of course, what the limit of the non-rescaled distributions $D_{N\alpha}(t)$ is. But, for each fixed $t > 0$, $D_{N\alpha}(t) \rightarrow 0$ as $N \rightarrow \infty$ by Theorem 1.2 and Proposition 3.5. Therefore, we need to replace $D_{N\alpha}(t)$ by $D_{N\alpha}(W_N(t))$ for an appropriate sequence $\{W_N(t)\}$ of positive functions in $t > 0$ which compensate for the flattening rate. Our sequence will satisfy the conditions $W_N(t) \rightarrow 0$ as $N \rightarrow \infty$ and

$$(84) \quad W_N(t) \rightarrow 0, \quad W_N(t)^{-1/N} \rightarrow W(t) \quad (N \rightarrow \infty),$$

for a function $W(t) > 1$. We then have the following limit distribution law:

THEOREM 4.3. — *Let $\{W_N\}$ satisfy (84). Then*

$$(85) \quad \lim_{N \rightarrow \infty} D_N(W_N(t)) = \text{Vol}_{M_P}(\{(\xi, \rho) \in \mathbb{C}^r \times (\mathbb{R}^*)^{m-r}; \Psi_\alpha(\xi, \rho) < \log W(t)\}).$$

In particular, by taking the function W_N as $W_N(t) = e^{-Nt}$, we have

$$(86) \quad \lim_{N \rightarrow \infty} D_N(e^{-Nt}) = \frac{1}{\pi^r} \int_{\{(\xi, \rho) \in \mathbb{C}^r \times (\mathbb{R}^*)^{m-r}; \Psi_\alpha^P(\xi, \rho) < t\}} L(\xi, \rho) \, dm(\xi) d\rho.$$

If x is in the interior P° of the polytope P , and if $\alpha_N \in NP \cap \mathbb{Z}^m$ be an approximate multiple of x , we have

$$(87) \quad \lim_{N \rightarrow \infty} D_N(e^{-Nt}) = \int_{\{\rho \in \mathbb{R}^m; b_x^P(\rho) < t\}} \det A(\rho) \, d\rho.$$

Proof. — We note that (86) follows from (85) and Lemma 3.6. So, first of all, we shall prove (85) for the boundary case. The set U_{v_0} introduced in Section 3 is dense in M_P , and hence we may consider the volume of the set

$$S_N(t) := \{(\xi, \zeta) \in \mathbb{C}^r \times (\mathbb{C}^*)^{m-r}; |\widehat{\varphi}_{N\alpha}^P(\xi, \zeta)|^2 > W_N(t)\},$$

so that $D_{N\alpha}(W_N(t)) = \text{Vol}_{M_P}(S_N(t))$. Let $(\xi, \zeta) \in S_N$. We write $\zeta = e^{\rho/2+i\theta}$. Then, by Proposition 3.5, we have

$$(88) \quad \Psi_\alpha(\xi, \rho) < \log \left(CN^{-(m+r)/2} W_N(t) (1 + a_N(\xi, \zeta)) \right)^{-1/N},$$

where C is a constant, and a_N is a function of order $O(N^{-1})$. Thus, we obtain

$$\begin{aligned} D_{N\alpha}(W_N(t)) &= \text{Vol}_{M_P} \left((\xi, \zeta) \in \mathbb{C}^r \times (\mathbb{C}^*)^{m-r}; \Psi_\alpha(\xi, \rho) \right. \\ &\quad \left. < \log \left(CN^{-(m+r)/2} W_N(t) (1 + a_N(\xi, \zeta)) \right)^{-\frac{1}{N}} \right). \end{aligned}$$

This combined with Lemma 3.6 shows the assertion.

Next, we shall show (87), or rather general version (85) for the case of interior points $x \in P^\circ$. Let $W_N(t)$ satisfy the condition (84). By Theorem 1.2, $|\varphi_{\alpha_N}(\rho)|^2 > W_N(t)$ implies

$$(89) \quad b_x^P(\rho) \leq \frac{1}{N} \log \left(CN^{m/2} (1 + a_N(\rho)) W_N(t)^{-1} \right) + \langle \rho - \rho_x^P, x_N \rangle,$$

where $C > 0$ is a constant and $a_N(\rho) = O(1/N)$ uniformly in $\rho \in \mathbb{R}^m$. From this and Lemma 3.9, the set

$$S_N(t) := \{\rho \in \mathbb{R}^m; |\varphi_{\alpha_N}(\rho)|^2 > W_N(t)\}$$

is bounded. Thus, since $x_N = \alpha_N/N - x = O(1/N)$, the function $\langle \rho - \rho_x^P, x_N \rangle$ is of order $O(1/N)$ uniformly around $S_N(t)$ for all N . The right hand side of (89) tends to $W(t)$ as $N \rightarrow \infty$ uniformly in ρ . Hence we have (85) for the case of interior points $x \in P^\circ$. □

Our final aim is to prove Theorem 1.3, which gives the asymptotic limit of the distribution functions $D_N(t)$ itself without any rescaling. Since, by Theorem 1.2 and Propositions 3.5 and 3.8, the monomial $|\widehat{\varphi}_{N\alpha}^P|^2$ decays exponentially away from the corresponding invariant torus $\mu_{P^{-1}}(\alpha)$, it is

obvious that $D_N(t)$ tends to zero as $N \rightarrow \infty$. So, a problem is to find the decay rate of $D_N(t)$ for any (but fixed) $t > 0$.

For every N and $0 \leq r \leq m$, we set

$$s_N = \log \left(\frac{N}{2\pi} \right)^{(m+r)/2} = \frac{m+r}{2} \log \left(\frac{N}{2\pi} \right), \quad r_N = \left(\frac{s_N}{N} \right)^{1/2}.$$

LEMMA 4.4. — *Let $t > 0$. Let $\alpha \in P$ be a lattice point with $\dim \mu_P^{-1}(\alpha) = m - r$ and lie in a face F with $\dim F = m - r$. Let $(\xi_N, \zeta_N) \in \mathbb{C}^r \times (\mathbb{C}^*)^{m-r}$ satisfy $|\widehat{\varphi}_{N\alpha}^P(\xi_N, \zeta_N)|^2 > t$. We write $\zeta_N = e^{(\rho_\alpha^F + u_N)/2 + i\theta_N}$. Then, we have*

$$|\xi_N|^2 + |u_N|^2 = O(r_N^2) = O(N^{-1} \log N)$$

locally uniformly in $t > 0$.

Proof. — For every $c > 0$, we set

$$(90) \quad B_\alpha(c) = \{(\xi, e^{(\rho_\alpha^F + u)/2} + i\theta) \in \mathbb{C}^r \times (\mathbb{C}^*)^{m-r}; |\xi|^2 + |u|^2 < c^2\}.$$

Then, by the argument in the proof of Proposition 3.5, we can find positive constants c_0, C_0 such that

$$c_0(|\xi|^2 + |u|^2) \leq \Psi_\alpha(\xi, \rho_\alpha^F + u) \leq C_0(|\xi|^2 + |u|^2) \text{ on } B_\alpha(c).$$

Therefore, by Proposition 3.5, we have

$$t \leq C e^{s_N - N\Psi_\alpha(\xi_N, \rho_\alpha^F + u_N)} \leq C e^{s_N - c_0 N(|\xi_N|^2 + |u_N|^2)}$$

with some constant C . Therefore, we obtain

$$|\xi_N|^2 + |u_N|^2 \leq \frac{C}{N} (\log(C/t) + s_N),$$

which implies the assertion. □

Thus, to obtain the asymptotic estimate of the distribution function $D_N(t)$, we need to find that of the monomial $|\widehat{\varphi}_{N\alpha}^P|^2$ on the ball $B_\alpha(cr_N)$ of radius $O(r_N)$ around the point $(0, \rho_\alpha^F)$, where $B_\alpha(cr_N)$ is defined in (90).

LEMMA 4.5. — *Let $c > 0$. We denote points $B_\alpha(cr_N)$ as $(\xi, \zeta) = (r_N w, e^{(\rho_\alpha^F + r_N u)/2 + i\theta})$ with $|w|^2 + |u|^2 \leq c^2$. Then we have*

$$\begin{aligned}
 & |\widehat{\varphi}_{N\alpha}^P(r_N w, \rho_\alpha^F + r_N u)|^2 \\
 (91) \quad & = c(P, \alpha) e^{s_N - s_N \langle H\Psi_\alpha(0, \rho_\alpha^F)(w, u), (w, u) \rangle / 2} (1 + O(N^{-1/2}(\log N)^{3/2})),
 \end{aligned}$$

where the Hessian $H\Psi_\alpha(0, \rho_\alpha^F)$ of the function Ψ_α on $\mathbb{C}^r \times \mathbb{R}^{m-r}$ at the point $(0, \rho_\alpha^F)$ is given by (62)

Proof. — By a Taylor expansion, we have

$$\Psi_\alpha(r_N w, \rho_\alpha^F + r_N u) = \frac{r_N^2}{2} \langle H\Psi_\alpha(0, \rho_\alpha^F)(w, u), (w, u) \rangle + R(r_N(w, u))$$

for $|w|^2 + |u|^2 \leq c^2$, where $R(r_N(w, u)) = O(r_N^3)$. In particular, we have

$$\begin{aligned}
 e^{-N\Psi_\alpha(r_N w, \rho_\alpha^F + r_N u)} & = e^{-s_N \langle H\Psi_\alpha(0, \rho_\alpha^F)(w, u), (w, u) \rangle / 2} (1 + O(Nr_N^3)) \\
 & = e^{-s_N \langle H\Psi_\alpha(0, \rho_\alpha^F)(w, u), (w, u) \rangle / 2} (1 + O(N^{-1/2}(\log N)^{3/2})).
 \end{aligned}$$

From this and the estimate in Proposition 3.5, the assertion follows for $1 \leq r \leq m - 1$. For $r = 0, m$, precisely the same argument replacing Ψ_α by b_α^P or $\log K$ with Theorem 1.2 and Proposition 3.8 will show the assertion. □

Remark. — Lemmas 4.4 and 4.5 are valid for interior points $x \in P^\circ$ and approximate multiples α_N . To see this, we note that the set

$$S_N(t) = \{\rho \in \mathbb{R}^m ; |\varphi_{\alpha_N}^P(\rho)|^2 > t\}$$

is contained in a ball around $\rho_x^P \in \mathbb{R}^m$. Thus, by the Taylor expansion for the function b_x^P , there exists constants $c > 0$ and $C > 0$ such that

$$b_x^P(\rho_x^P + u) - \langle u, x_N \rangle \geq c|u|^2, \quad |u| \leq C.$$

From this, we have $|u| = O(r_N)$ if $\rho_x^P + u \in S_N(t)$, where $r_N = (s_N/N)^{1/2}$ and $s_N = \log \left(\frac{N}{2\pi}\right)^{m/2}$. By Theorem 1.2 and the Taylor expansion, we have

$$\begin{aligned}
 |\varphi_{\alpha_N}^P(\rho_x^P + r_N u)|^2 & = c(P, x) e^{s_N - s_N \langle A(P, x)u, u \rangle / 2} (1 + O(N^{-1/2}(\log N)^{-3/2})), \\
 & \quad |u| \leq C.
 \end{aligned}$$

Proof of Theorem 1.3. — For every $t > 0$, we set

$$S_N(t) = \{\eta = (\xi, \zeta) \in \mathbb{C}^r \times (\mathbb{C}^*)^{m-r}; |\widehat{\varphi}_{N\alpha}^P(\xi, \zeta)|^2 > t\}.$$

Then, by Lemma 4.4, there is a constant $c > 0$ (depending on a fixed $t > 0$) such that $S_N(t) \subset B_\alpha(cr_N)$. Let $(\xi, \zeta) = (r_N w, e^{\rho_\alpha^F + r_N u})/2 + i\theta) \in B_\alpha(cr_N)$. Then, by Lemma 4.5, $(\xi, \zeta) \in S_N(t)$ if and only if

$$t < c(P, \alpha) e^{s_N - s_N \langle H\Psi_\alpha(0, \rho_\alpha^F)(w, u), (w, u) \rangle / 2} (1 + a_N(w, u)),$$

where $a_N(w, u)$ is a function of order $N^{-1/2}(\log N)^{3/2}$ uniformly in (w, u) with $|w|^2 + |u|^2 \leq c^2$, and $c(P, \alpha)$ is define in (73). This is equivalent to the following estimate:

$$\langle H\Psi_\alpha(w, u), (w, u) \rangle / 2 < 1 + \frac{1}{s_N} \log \left(\frac{c(P, \alpha)}{t} (1 + a_N(w, u)) \right).$$

Therefore, by Lemma 3.6, we have

$$\begin{aligned} D_{N\alpha}(t) &= \frac{1}{\pi^r} \int_{\{(\xi, \rho) = (r_N w, \rho_\alpha^F + r_N u); \langle H(w, u), (w, u) \rangle / 2 < 1 + \frac{1}{s_N} \log(c(P, \alpha)(1 + a(w, u))/t)\}} L(\xi, \rho) dm(\xi) d\rho, \end{aligned}$$

where we set, for simplicity, $H = H\Psi_\alpha(0, \rho_\alpha^F)$. Changing the variables (ξ, ρ) to (w, u) , we have

$$\begin{aligned} D_{N\alpha}(t) &= \frac{r_N^{m+r}}{\pi^r} \int_{\{(w, u); \langle H(w, u), (w, u) \rangle / 2 < 1 + \frac{1}{s_N} \log(c(P, \alpha)(1 + a_N(w, u))/t)\}} L(r_N w, \rho_\alpha^F + r_N u) dm(w) du. \end{aligned}$$

A Taylor expansion gives

$$L(r_N w, \rho_\alpha^F + r_N u) = L(0, \rho_\alpha^F) + R_N(w, u)$$

with the error term $R_N(w, u) = O(r_N)$ uniformly for $|w|^2 + |u|^2 \leq c^2$. Thus, we obtain

$$\begin{aligned} r_N^{-(m+r)} D_{N\alpha}(t) &= \frac{L(0, \rho_\alpha^F)}{\pi^r} \int_{\{(w, u); \langle H(w, u), (w, u) \rangle / 2 < 1 + \frac{1}{s_N} \log(c(P, \alpha)(1 + a_N(w, u))/t)\}} dm(w) du + O(r_N). \end{aligned}$$

Note that $s_N \rightarrow \infty$ while $r_N \rightarrow 0$ as $N \rightarrow \infty$. The function $a_N(w, u)$ tends to zero uniformly in $|w|^2 + |u|^2 \leq c^2$. Therefore, we conclude, for a fixed $t > 0$,

$$\begin{aligned} \lim_{N \rightarrow \infty} r_N^{-(m+r)} D_{N\alpha}(t) &= \frac{L(0, \rho_\alpha^F)}{\pi^r} \int_{\{(w,u) \in \mathbb{C}^r \times \mathbb{R}^{m-r}; \langle H(w,u), (w,u) \rangle / 2 < 1\}} dm(w) du. \\ &= \frac{2^{(m+r)/2}}{\pi^r} \frac{L(0, \rho_\alpha^F)}{\sqrt{\det H}} \text{Vol}_{M_P}(x \in \mathbb{R}^{m+r}; |x| < 1). \end{aligned}$$

By (64) and the well-known formula for the volume of the unit disk in \mathbb{R}^{m+r} , we conclude the assertion. For the case of the interior points of the polytope P , these can be proved by using the fact proved in Remark given above. \square

BIBLIOGRAPHY

- [Be] M.V. BERRY, Regular and irregular semiclassical wavefunctions, *J. Phys. A* 10 (1977), 2083-2091.
- [BHO] M.V. BERRY, J. H. HANNAY, and A.M. OZORIO de ALMEIDA, Intensity moments of semiclassical wavefunctions, *J. Phys. D* 8 (1983), 229-242.
- [De] T. DELZANT, Hamiltoniens périodiques et image convexe de l'application moment, *Bull. Soc. Math. France* 116 (1988), 315-339.
- [FE] V. I. FALCKO and K. B. EFETOV, Statistics of wave functions in mesoscopic systems, *J. Math. Phys.* 37 (1996), 4935-4967.
- [Fu] W. FULTON, Introduction to Toric Varieties, *Annals of Math. Study* 131, Princeton Univ. Press, Princeton, 1993.
- [GKZ] I. M. GELFAND, M. M. KAPRANOV and A. V. ZELEVINSKY, Discriminants, resultants, and multidimensional determinants, *Mathematics: Theory and Applications*, Birkhäuser, Boston, 1994.
- [Gu] V. GUILLEMIN, Moment Maps and Combinatorial Invariants of Hamiltonian T^n -Spaces, *Progress in Math.* 122, Birkhäuser, Boston, 1994.
- [He] D. A. HEJHAL, On eigenfunctions of the Laplacian for Hecke triangle groups, *Emerging applications of number theory* (Minneapolis, MN, 1996), *IMA Vol. Math. Appl.* 109, Springer-Verlag, New York, 1999, pp. 291-315.
- [HR] D. A. HEJHAL and B. N. RACKNER, On the topography of Maass waveforms for $\text{PSL}(2, Z)$, *Experiment. Math.* 1 (1992), 275-305.
- [Hö] L. HÖRMANDER, *The Analysis of Linear Partial Differential Operators, I* (Second Ed.), Springer-Verlag, New York, 1990.
- [Ka] N. M. KATZ, Sato-Tate equidistribution of Kurlberg-Rudnick sums, *Internat. Math. Res. Notices* (2001), 711-728.
- [KR] P. KURLBERG and Z. RUDNICK, Value distribution for eigenfunctions of desymmetrized quantum maps, *Internat. Math. Res. Notices* (2001), 985-1002.

- [LS] E. LERMAN and N. SHIROKOVA, Completely integrable torus actions on symplectic cones, *Math. Res. Lett.* 9 (2002), 105-115.
- [Mi] A. D. MIRLIN, Statistics of energy levels and eigenfunctions in disordered systems, *Phys. Rep.* 326 (2000), 259-382.
- [MF] A. D. MIRLIN and Y. V. FYODOROV, Distribution of local densities of states, order parameter function, and critical behavior near the Anderson transition, *Phys. Rev. Lett.* 72, (1994), 526-529.
- [PA] V. N. PRIGODIN and B. L. ALTSHULER, Long-range spatial correlations of eigenfunctions in quantum disordered systems, *Phys. Rev. Lett.* 80 (1998), 1944-1947.
- [STZ] B. SHIFFMAN, T. TATE and S. ZELDITCH, Harmonic analysis on toric varieties, *Explorations in Complex and Riemannian Geometry*, Contemporary Math., 332, Amer. Math. Soc., Providence, RI, pp. 267-286.
- [SZ1] B. SHIFFMAN and S. ZELDITCH, Distribution of zeros of random and quantum chaotic sections of positive line bundles, *Comm. Math. Phys.* 200 (1999), 661-683.
- [SZ2] B. SHIFFMAN and S. ZELDITCH, Random polynomials with prescribed Newton polytope, *J. Amer. Math. Soc.* 17 (2004), 49-108.
- [SS] M. SREDNICKI and F. STIERNELOF, Gaussian fluctuations in chaotic eigenstates, *J. Phys. A* 29 (1996), 5817-5826.
- [TZ] J. TOTH and S. ZELDITCH, L^p -norms of eigenfunctions in the completely integrable case, *Ann. Henri Poincaré* 4 (2003), 343-368.
- [Y] S.-T. YAU, Open problems in geometry, *Differential Geometry: Partial Differential Equations on Manifolds* (Los Angeles, CA, 1990), Proc. Sympos. Pure Math., 54, Part 1, Amer. Math. Soc., Providence, RI, 1993, pp. 1-28.
- [Ze] S. ZELDITCH, Szegő kernels and a theorem of Tian, *Internat. Math. Res. Notices* (1998), 317-331.

Bernard SHIFFMAN, Steve ZELDITCH,
 Johns Hopkins University
 Department of Mathematics
 Baltimore, MD 21218 (USA).

shiffman@math.jhu.edu
 zelditch@math.jhu.edu

Tatsuya TATE,
 Department of Mathematics
 Keio University
 Keio University 3-14-1 Hiyoshi Kohoku-ku
 Yokohama, 223-8522 (Japan).

tate@math.keio.ac.jp