



ANNALES

DE

L'INSTITUT FOURIER

Marcia EDSON & Luca Q. ZAMBONI

On the Number of Partitions of an Integer in the m -bonacci Base

Tome 56, n° 7 (2006), p. 2271-2283.

http://aif.cedram.org/item?id=AIF_2006__56_7_2271_0

© Association des Annales de l'institut Fourier, 2006, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

ON THE NUMBER OF PARTITIONS OF AN INTEGER IN THE m -BONACCI BASE

by Marcia EDSON & Luca Q. ZAMBONI

ABSTRACT. — For each $m \geq 2$, we consider the m -bonacci numbers defined by $F_k = 2^k$ for $0 \leq k \leq m-1$ and $F_k = F_{k-1} + F_{k-2} + \cdots + F_{k-m}$ for $k \geq m$. When $m = 2$, these are the usual Fibonacci numbers. Every positive integer n may be expressed as a sum of distinct m -bonacci numbers in one or more different ways. Let $R_m(n)$ be the number of partitions of n as a sum of distinct m -bonacci numbers. Using a theorem of Fine and Wilf, we obtain a formula for $R_m(n)$ involving sums of binomial coefficients modulo 2. In addition we show that this formula may be used to determine the number of partitions of n in more general numeration systems including generalized Ostrowski number systems in connection with Episturmian words.

RÉSUMÉ. — Pour $m \geq 2$, on définit les nombres de m -bonacci $F_k = 2^k$ pour $0 \leq k \leq m-1$ et $F_k = F_{k-1} + F_{k-2} + \cdots + F_{k-m}$ pour $k \geq m$. Dans le cas $m = 2$, on retrouve les nombres de Fibonacci. Chaque entier positif n s'écrit comme une somme distincte de nombres de m -bonacci d'une ou plusieurs façons. Soit $R_m(n)$ le nombre de partitions de n en base m -bonacci. En utilisant un théorème de Fine et Wilf on déduit une formule pour $R_m(n)$ comme somme de coefficients binomiaux modulo 2. De plus, nous montrons que cette formule peut-être utilisée pour déterminer le nombre de partitions de n dans des systèmes généraux de numération incluant les systèmes de nombres d'Ostrowski généralisés associés aux suites episturmiennes.

1. Introduction and Preliminaries

For each $m \geq 2$, we define the m -bonacci numbers by $F_k = 2^k$ for $0 \leq k \leq m-1$ and $F_k = F_{k-1} + F_{k-2} + \cdots + F_{k-m}$ for $k \geq m$. When $m = 2$, these are the usual Fibonacci numbers. We denote by $\{0, 1\}^*$ the set of all words $w = w_1 w_2 \cdots w_k$ with $w_i \in \{0, 1\}$. Each positive integer n may be expressed as a sum of distinct m -bonacci in one or more different ways.

Keywords: Numeration systems, Fibonacci numbers, Fine and Wilf theorem, episturmian words.

Math. classification: 11B39, 11B50, 68R15.

That is we can write $n = \sum_{i=1}^k w_i F_{k-i}$ where $w_i \in \{0, 1\}$ and $w_1 = 1$. We call the associated $\{0, 1\}$ -word $w_1 w_2 \cdots w_k$ a *representation* of n . One way of obtaining such a representation is by applying the “greedy algorithm”. This gives rise to a representation of n of the form $w = w_1 w_2 \cdots w_k$ with the property that w does not contain m consecutive 1’s. Such a representation of n is necessarily unique and is called the *m -Zeckendorff representation* of n , denoted $Z_m(n)$ [13]. For example, taking $m = 2$ and applying the greedy algorithm to $n = 50$ we obtain $50 = 34 + 13 + 3 = F_7 + F_5 + F_2$ which gives rise to the representation $Z_2(50) = 10100100$. A $\{0, 1\}$ -word w beginning in 1 and having no occurrences of 1^m will be called a *m -Zeckendorff word*.

Other representations arise from the fact that an occurrence of 10^m in a given representation of n may be replaced by 01^m to obtain another representation of n , and conversely. Thus a number n has a unique representation in the m -bonacci base if and only if $Z_m(n)$ does not contain any occurrences of 0^m . For example, again taking $m = 2$ and $n = 50$ we obtain the following 6 representations (arranged in decreasing lexicographic order):

10100100
10100011
10011100
10011011
1111100
1111011

We are interested in the sequence $R_m(n)$ which counts the number of distinct partitions of n in the m -bonacci base. More precisely, given $n \in \mathbb{Z}^{>0}$ we set

$$\Omega_m(n) = \{w = w_1 w_2 \cdots w_k \in \{0, 1\}^* \mid w_1 = 1 \text{ and } n = \sum_{i=1}^k w_i F_{k-i}\}$$

and put $R_m(n) = \#\Omega_m(n)$. For $w \in \Omega_m(n)$ we will sometimes write $R_m(w)$ for $R_m(n)$. Also we let $R_m^{\leq}(w)$ denote the number of representations of n which are less or equal to w in the lexicographic order. As $Z_m(n)$ is the largest representation of n with respect to the lexicographic order, it follows that $R_m(n) = R_m^{\leq}(Z_m(n))$.

In a 1968 paper L. Carlitz [3] studied the multiplicities of representations of n as sums of distinct Fibonacci numbers; he obtained recurrence relations for $R_2(n)$ and explicit formulae for $R_2(n)$ in the case $Z_2(n)$ contains 1, 2 or 3 Fibonacci numbers. He states in the paper however that a general formula for the number of partitions of n in the Fibonacci base appears

to be very complicated. In [1] J. Berstel derives a formula for $R_2(n)$ as a product of 2×2 matrices (see Proposition 4.1 in [1]). Recently, P. Kocábová, Z. Masáková, and E. Pelantová [10] extended Berstel's result to $R_m(n)$ for all $m \geq 2$ again as a product of 2×2 matrices.

In this paper we give a formula for $R_m(n)$ involving sums of binomial coefficients modulo 2. Our proof makes use of the well known Fine and Wilf Theorem [4]. In order to state our main result, we first consider a special factorization of $Z_m(n)$: Either $Z_m(n)$ contains no occurrences of 0^m (in which case $R_m(n) = 1$), or $Z_m(n)$ can be factored uniquely in the form

$$Z_m(n) = V_1 U_1 V_2 U_2 \cdots V_N U_N W$$

where

- V_1, V_2, \dots, V_N and W do not contain any occurrences of 0^m .
- 0^{m-1} is not a suffix of V_1, V_2, \dots, V_N .
- Each U_i is of the form

$$U_i = 10^{m-1} x_k 0^{m-1} x_{k-1} \cdots 0^{m-1} x_0 0^m$$

with $x_i \in \{0, 1\}$.

We shall refer to this factorization as the *principal factorization* of $Z_m(n)$ and call the U_i *indecomposable factors*. We observe that in the special case of $m = 2$, the factors V_i are empty. Each indecomposable factor U_i may be coded by a positive integer r_i whose base 2 expansion is $1x_k x_{k-1} \cdots x_0$, in other words $r_i = 1 \cdot 2^{k+1} + x_k \cdot 2^k + \cdots + x_1 \cdot 2 + x_0$.

Given a positive integer r whose base 2 expansion is $1x_k x_{k-1} \cdots x_0$, we set

$$[r] = 10^{m-1} x_k 0^{m-1} x_{k-1} \cdots 0^{m-1} x_0 0^m.$$

We now state our main result:

THEOREM 1.1. — *Let $m \geq 2$. Given a positive integer n , let $Z_m(n) = V_1 U_1 V_2 U_2 \cdots V_N U_N W$ be the principal factorization of the m -Zeckendorff representation of n as defined above. Then the number of distinct partitions of n as sums of distinct m -bonacci numbers is given by*

$$R_m(n) = \prod_{i=1}^N \sum_{j=0}^{r_i} \binom{2r_i - j}{j} \pmod{2}$$

where $[r_i] = U_i$ for each $1 \leq i \leq N$.

2. Proof of Theorem 1.1

Let $Z_m(n) = V_1U_1V_2U_2 \cdots V_NU_NW$ be the principal factorization of $Z_m(n)$ described above. Then the number of partitions of n is simply the product of the number of partitions of each indecomposable factor:

$$(2.1) \quad R_m(n) = \prod_{i=1}^N R_m(U_i).$$

In fact, any representation of n as a sum of distinct m -bonacci numbers may be factored in the form

$$V_1U'_1V_2U'_2 \cdots V_NU'_NW$$

where for each $1 \leq i \leq N$, U'_i is an equivalent representation of U_i . To see this we first observe that since the V_i and W contain no 0^m , we have $R_m(V_i) = R_m(W) = 1$. So the only way that V_i or W could change in an alternate representation of n would be as a result of a neighboring indecomposable factor. If V_i contains an occurrence of 1, then since V_i does not end in 0^{m-1} the last occurrence of 1 in V_i can never be followed by 0^m . In other words the last 1 in V_i can never move into the U_i that follows. If V_i contains no occurrences of 1, then $V_i = 0^r$ with $r < m - 1$. Since the indecomposable factor U_{i-1} preceding V_i ends in Km many consecutive 0's (for some $K \geq 1$), any equivalent representation of U_{i-1} either ends in 0^m or in 1^m , and since V_i does not begin in 0^m , any representation of U_{i-1} terminating in 1^m will never be followed by 0^m . In other words, no 1 in U_{i-1} can ever move into V_i or in the following U_i . A similar argument applies to the indecomposable factor U_N preceding W .

Thus in view of (2.1) above, in order to prove Theorem 1.1, it remains to show that for each positive integer $r = 1 \cdot 2^{k+1} + x_k \cdot 2^k + \cdots + x_1 \cdot 2 + x_0$, we have

$$(2.2) \quad R_m([r]) = \sum_{j=0}^r \binom{2r-j}{j} \pmod{2}.$$

For each positive integer n there is a natural decomposition of the set $\Omega_m(n)$ of all partitions of n in the m -bonacci base: Let F be the largest m -bonacci number less or equal to n . We denote by $\Omega_m^+(n)$ the set of all partitions of n involving F and $\Omega_m^-(n)$ the set of all partitions of n not involving F , and set $R_m^+(n) = \#\Omega_m^+(n)$ and $R_m^-(n) = \#\Omega_m^-(n)$. Clearly

$$R_m(n) = R_m^+(n) + R_m^-(n).$$

We will make use of the following recursive relations:

LEMMA 2.1. — Let $U = 10^{m-1}x_k0^{m-1}x_{k-1} \cdots 0^{m-1}x_00^m$ with $x_i \in \{0, 1\}$. Then

$$R_m^+(10^{m-1}10^{m-1}x_k0^{m-1}x_{k-1} \cdots 0^{m-1}x_00^m) = R_m(U) = R_m^+(U) + R_m^-(U)$$

$$R_m^-(10^{m-1}10^{m-1}x_k0^{m-1}x_{k-1} \cdots 0^{m-1}x_00^m) = R_m^-(U)$$

$$R_m^+(10^{m-1}00^{m-1}x_k0^{m-1}x_{k-1} \cdots 0^{m-1}x_00^m) = R_m^+(U)$$

$$R_m^-(10^{m-1}00^{m-1}x_k0^{m-1}x_{k-1} \cdots 0^{m-1}x_00^m) = R_m(U) = R_m^+(U) + R_m^-(U)$$

Proof. — It is easy to see that $w \in \Omega_m^+(10^{m-1}U)$ if and only if w is of the form $w = 10^{m-1}w'$ for some $w' \in \Omega_m(U)$. Whence $R_m^+(10^{m-1}U) = R_m(U)$. Similarly, $w \in \Omega_m^-(10^{m-1}U)$ if and only if w is of the form $w = 01^mw'$ for some $w' \in \Omega_m^-(U)$. Whence $R_m^-(10^{m-1}U) = R_m^-(U)$. A similar argument applies to the remaining two identities. \square

Fix a positive integer $r = 1 \cdot 2^{k+1} + x_k \cdot 2^k + \cdots + x_1 \cdot 2 + x_0$. The above lemma can be used to compute $R_m([r])$ as follows: We construct a tower of $k + 2$ levels L_0, L_1, \dots, L_{k+1} , where each level L_i consists of an ordered pair (a, b) of positive integers. We start with level 0 by setting $L_0 = (1, 1)$. Then L_{i+1} is obtained from L_i according to the value of x_i . If $L_i = (a, b)$, then $L_{i+1} = (a, a + b)$ if $x_i = 0$, and $L_{i+1} = (a + b, b)$ if $x_i = 1$. It follows from the above Lemma that $L_{k+1} = (R_m^+([r]), R_m^-([r]))$. Hence $R_m([r])$ is the sum of the entries of level L_{k+1} .

By the well known Fine and Wilf Theorem [4], given a pair of relatively prime numbers (p, q) , there exists a $\{0, 1\}$ -word w of length $p + q - 2$ (unique up to isomorphism) having periods p and q , and if p and q are both greater than 1, then this word contains both 0's and 1's; in other words $1 = \gcd(p, q)$ is not a period. We call such a word a *Fine and Wilf word* relative to (p, q) . Moreover it can be shown (see [12] for example) that if both p and q are greater than 1, then the suffixes of w of lengths p and q begin in different symbols. We denote by $FW(p, q)$ the unique Fine and Wilf word relative to (p, q) with the property that its suffix of length p begins in 0 and its suffix of length q begins in 1.

We now apply this to the ordered pair $(p, q) = (R_m^+([r]), R_m^-([r]))$. It is well known that $FW(R_m^+([r]), R_m^-([r]))01$ is given explicitly by the following composition of morphisms:

$$FW(R_m^+([r]), R_m^-([r]))01 = \tau_{x_0} \circ \tau_{x_1} \circ \cdots \circ \tau_{x_k}(01)$$

where

$$\begin{aligned} \tau_0(0) &= 0 & \tau_0(1) &= 01 \\ \tau_1(0) &= 10 & \tau_1(1) &= 1 \end{aligned}$$

(see for instance [5, 12]).

Let

$$\alpha(r) = |FW(R_m^+([r]), R_m^-([r]))01|_1$$

and

$$\beta(r) = |FW(R_m^+([r]), R_m^-([r]))01|_0$$

in other words, $\alpha(r)$ is the number of occurrences of 1 in

$$FW(R_m^+([r]), R_m^-([r]))01$$

and $\beta(r)$ the number of 0's in

$$FW(R_m^+([r]), R_m^-([r]))01.$$

In summary:

$$\begin{aligned} R_m([r]) &= R_m^+([r]) + R_m^-([r]) \\ &= R_m^+([r]) + R_m^-([r]) - 2 + 2 \\ &= |FW(R_m^+([r]), R_m^-([r]))| + 2 \\ &= |FW(R_m^+([r]), R_m^-([r]))01| \\ &= |\tau_{x_0} \circ \tau_{x_1} \circ \dots \circ \tau_{x_k}(01)| \\ &= |\tau_{x_0} \circ \tau_{x_1} \circ \dots \circ \tau_{x_k}(01)|_1 + |\tau_{x_0} \circ \tau_{x_1} \circ \dots \circ \tau_{x_k}(01)|_0 \\ &= \alpha(r) + \beta(r) \\ &= |\tau_1 \circ \tau_{x_0} \circ \tau_{x_1} \circ \dots \circ \tau_{x_k}(01)|_1 \\ &= \alpha(2r + 1). \end{aligned}$$

The key step in the proof of Theorem 1.1 is to replace above the sum of the periods $R_m^+([r]) + R_m^-([r])$ of the Fine and Wilf word $FW(R_m^+([r]), R_m^-([r]))$ by the sum of the number of occurrences of 0's and 1's in $FW(R_m^+([r]), R_m^-([r]))01$. The following basic identities are readily verified:

- $\alpha(1) = \beta(1) = 1.$
- $\alpha(2r) = \alpha(r).$
- $\beta(2r) = \alpha(r) + \beta(r).$
- $\alpha(2r + 1) = \alpha(r) + \beta(r).$
- $\beta(2r + 1) = \beta(r).$
- $\beta(r) = \alpha(r + 1).$

Summarizing we have

PROPOSITION 2.2. — *Let $U = 10^{m-1}x_k0^{m-1}x_{k-1}\dots0^{m-1}x_00^m$ with $x_i \in \{0, 1\}$. Let r be the number whose base 2 expansion is given by $1x_kx_{k-1}\dots x_0$. Then $R_m(U) = \alpha(2r + 1)$ where the sequence $\alpha(r)$ is defined recursively by:*

- $\alpha(1) = 1$
- $\alpha(2r) = \alpha(r)$
- $\alpha(2r + 1) = \alpha(r) + \alpha(r + 1)$.

We now consider a new function $\psi(r)$ defined by $\psi(1) = 1$, and for $r \geq 1$

$$\psi(r + 1) = \sum_{j=0}^{2j \leq r} \binom{r-j}{j} \pmod{2}.$$

We will show that $\psi(r)$ and $\alpha(r)$ satisfy the same recursive relations, namely: $\psi(2r) = \psi(r)$ and $\psi(2r + 1) = \psi(r) + \psi(r + 1)$. Thus $\alpha(r) = \psi(r)$ for each r thereby establishing formula (2.2).

We shall make use of the following lemma:

LEMMA 2.3. — $\binom{n}{k} \pmod{2} = \binom{2n+1}{2k} \pmod{2} + \binom{2n}{2k+1} \pmod{2}$.

Proof. — This follows immediately from the so-called Lucas' identities: $\binom{2n}{2k+1} = 0 \pmod{2}$ for $0 \leq k \leq n - 1$, and $\binom{n}{k} = \binom{2n+1}{2k} \pmod{2}$ for $0 \leq k \leq n$. \square

PROPOSITION 2.4. — For $r \geq 0$ we have $\psi(2r + 2) = \psi(r + 1)$ and for $r \geq 1$ we have $\psi(2r + 1) = \psi(r) + \psi(r + 1)$.

Proof. — By Lemma 2.3 we have

$$\begin{aligned} \psi(r + 1) &= \sum_{j=0}^{2j \leq r} \binom{r-j}{j} \pmod{2} \\ &= \sum_{j=0}^{2j \leq r} \left(\binom{2r-2j+1}{2j} \pmod{2} + \binom{2r-2j}{2j+1} \pmod{2} \right) \\ &= \sum_{i=0}^r \binom{2r+1-i}{i} \pmod{2} \\ &= \psi(2r + 2). \end{aligned}$$

As for the second recursive relation we have

$$\begin{aligned} \psi(2r + 1) &= \sum_{j=0}^r \binom{2r-j}{j} \pmod{2} \\ &= \sum_{i=0}^{2i \leq r} \binom{2r-2i}{2i} \pmod{2} + \sum_{i=0}^{2i \leq r-1} \binom{2r-2i-1}{2i+1} \pmod{2} \end{aligned}$$

But

$$\begin{aligned}
 \binom{2r-2i}{2i} \pmod{2} &= \frac{(2r-2i)!}{(2i)!(2r-4i)!} \pmod{2} \\
 &= \frac{(2r-2i+1)!}{(2i)!(2r-4i+1)!} \pmod{2} \\
 &= \binom{2r-2i+1}{2i} \pmod{2} \\
 &= \binom{r-i}{i} \pmod{2} \text{ by Lemma 2.3.}
 \end{aligned}$$

Hence

$$\sum_{i=0}^{2i \leq r} \binom{2r-2i}{2i} \pmod{2} = \sum_{i=0}^{2i \leq r} \binom{r-i}{i} \pmod{2} = \psi(r+1).$$

Similarly

$$\begin{aligned}
 \binom{2r-2i-1}{2i+1} \pmod{2} &= \frac{(2r-2i-1)!}{(2i+1)!(2r-4i-2)!} \pmod{2} \\
 &= \frac{(2r-2i-1)!}{(2i)!(2r-4i-1)!} \pmod{2} \\
 &= \binom{2r-2i-1}{2i} \pmod{2} \\
 &= \binom{r-1-i}{i} \pmod{2} \text{ by Lemma 2.3.}
 \end{aligned}$$

Hence

$$\sum_{i=0}^{2i \leq r-1} \binom{2r-2i-1}{2i+1} \pmod{2} = \sum_{i=0}^{2i \leq r-1} \binom{r-1-i}{i} \pmod{2} = \psi(r).$$

It follows that $\psi(2r+1) = \psi(r) + \psi(r+1)$. □

Having established that $\alpha(r) = \psi(r)$ for each $r \geq 1$, we deduce that:

COROLLARY 2.5. — *Let $U = 10^{m-1}x_k0^{m-1}x_{k-1} \cdots 0^{m-1}x_00^m$ with $x_i \in \{0, 1\}$. Let r be the number whose base 2 expansion is given by $1x_kx_{k-1} \cdots x_0$. Then $R_m(U) = \sum_{j=0}^r \binom{2r-j}{j} \pmod{2}$.*

This concludes our proof of Theorem 1.1.

3. Concluding Remarks

3.1. A formula for $R_m^{\leq}(w)$

Our proof applies more generally to give a formula for $R_m^{\leq}(w)$ for each representation w of n . In other words, given $w \in \Omega_m(n)$, then either w does not contain any occurrences of 0^m (in which case $R_m^{\leq}(w) = 1$) or w may be factored in the form

$$w = V_1 U_1 V_2 U_2 \cdots V_N U_N W$$

where the V_i and W do not contain any occurrences of 0^m and the V_i do not end in 0^{m-1} , and where the U_i are of the form

$$U_i = 10^{m-1} x_k 0^{m-1} x_{k-1} \cdots 0^{m-1} x_0 0^m$$

with $x_i \in \{0, 1\}$. Each factor U_i is coded by a positive integer r_i whose base 2 expansion is $1x_k x_{k-1} \cdots x_0$. It is easy to see that any representation of n less or equal to w may be factored in the form

$$V_1 U'_1 V_2 U'_2 \cdots V_N U'_N W$$

where for each $1 \leq i \leq N$, U'_i is an equivalent representation of U_i . Hence $R_m^{\leq}(w) = \prod_{i=1}^N R_m(U_i)$ from which it follows that

$$R_m^{\leq}(w) = \prod_{i=1}^N \sum_{j=0}^{r_i} \binom{2r_i - j}{j} \pmod{2}.$$

3.2. Episturmian numeration systems

Let A be a finite non-empty set. Associated to an infinite word $\omega = \omega_1 \omega_2 \omega_3 \dots \in A^{\mathbb{N}}$ is a non-decreasing sequence of positive integers $\mathcal{E}(\omega) = E_1, E_2, E_3, \dots$ defined recursively as follows: $E_1 = 1$, and for $k \geq 1$, the quantity E_{k+1} is defined by the following rule: If $\omega_{k+1} \neq \omega_j$ for each $1 \leq j \leq k$, then set

$$E_{k+1} = 1 + \sum_{j=1}^k E_j.$$

Otherwise let $\ell \leq k$ be the largest integer such that $\omega_{k+1} = \omega_\ell$, and set

$$E_{k+1} = \sum_{j=\ell}^k E_j.$$

In particular we note that $E_{k+1} = E_k$ if and only if $\omega_{k+1} = \omega_k$.

Set $\mathcal{N}(\omega) = \{E_k | k \geq 1\}$. For $E \in \mathcal{N}(\omega)$ let $k \geq 1$ be such that $E = E_k$. We define $\sigma(E) = \omega_k$ and say that E is *based* at $\omega_k \in A$. We also define the quantity $\rho(E)$, which we call the *multiplicity* of E , by

$$\rho(E) = \#\{i \geq 1 | E = E_i\}.$$

We can write $\mathcal{N}(\omega) = \{x_1, x_2, x_3, \dots\}$ where for each $i \geq 1$ we have $x_i < x_{i+1}$. Thus we have that $\omega = \sigma(x_1)^{\rho(x_1)}\sigma(x_2)^{\rho(x_2)} \dots$

It can be verified that the set $\mathcal{N}(\omega)$ defines a numeration system (see [8]). More precisely, each positive integer n may be written as a sum of the form

$$(3.1) \quad n = m_k x_k + m_{k-1} x_{k-1} + \dots + m_1 x_1$$

where for each $1 \leq i \leq k$ we have $0 \leq m_i \leq \rho(x_i)$ and $m_k \geq 1$. While such a representation of n is not necessarily unique, one way of obtaining such a representation is to use the “greedy algorithm”. In this case we call the resulting representation the Zeckendorff representation of n and denote it $Z_\omega(n)$. We call the above numeration system a *generalized Ostrowski system* or an *Episturmian numeration system*. In fact, the quantities E_i are closely linked to the lengths of the palindromic prefixes of the characteristic Episturmian word associated to the directive sequence ω (see [6, 7, 8, 9]). In case $\#A = 2$, this is known as the Ostrowski numeration system (see [1, 2, 11]). In case $A = \{1, 2, \dots, m\}$ and ω is the periodic sequence $\omega = (1, 2, 3, \dots, m,)^\infty$, then the resulting numeration system is the m -bonacci system defined earlier.

Given an infinite word $\omega = \omega_1\omega_2\omega_3 \dots \in A^\mathbb{N}$, we are interested in the number of distinct ways of writing each positive integer n as a sum of the form (3.1). More precisely, denoting by \hat{A} the set $\{\hat{a} | a \in A\}$, we set $R_\omega(n) = \#\Omega_\omega(n)$ where $\Omega_\omega(n)$ is the set of all expressions of the form

$$(3.2) \quad \widehat{\sigma(x_k)}^{m_k} \sigma(x_k)^{\rho(x_k)-m_k} \widehat{\sigma(x_{k-1})}^{m_{k-1}} \dots \widehat{\sigma(x_1)}^{m_1} \sigma(x_1)^{\rho(x_1)-m_1}$$

in $(A \cup \hat{A})^*$, such that $n = m_k x_k + m_{k-1} x_{k-1} + \dots + m_1 x_1$ where $\mathcal{N}(\omega) = \{x_1, x_2, x_3, \dots | 1 = x_1 < x_2 < x_3 \dots\}$ and where $0 \leq m_i \leq \rho(x_i)$ and $m_k \geq 1$.⁽¹⁾ For $w \in \Omega_\omega(n)$ we sometimes write $R_\omega(w)$ for $R_\omega(n)$.

Just as in the previous section, we begin with a unique special factorization of the Zeckendorff representation of n . In this case, this factorization

⁽¹⁾ Our notation here differs somewhat from that of Justin and Pirillo in [8]. For instance, in [8] the authors use the notation \tilde{a} for in lieu of our \hat{a} . Also instead of the expression (3.2), they consider the reverse of this word.

was originally defined by Justin and Pirillo (see Theorem 2.7 in [8]):

$$Z_\omega(n) = V_1 U_1 V_2 U_2 \cdots V_N U_N W$$

where for each $1 \leq i \leq N$ we have that U_i is a a_i -based maximal semigood multiblock for some $a_i \in A$. Moreover any other representation of n may be factored in the form

$$Z_\omega(n) = V_1 U'_1 V_2 U'_2 \cdots V_N U'_N W$$

where U'_i is an equivalent representation of U_i (see Theorem 2.7 in [8]). Thus as before (see (2.1)) we have

$$R_\omega(n) = \prod_{i=1}^N R_\omega(U_i).$$

For each $1 \leq i \leq N$ the factor U_i corresponds to a sum of the form

$$m_K x_K + m_{K-1} x_{K-1} + \cdots + m_k x_k$$

for some $K > k$ with $m_K \neq 0$, and for each $K \geq j \geq k$ we have that if $m_j \neq 0$, then $\sigma(x_j) = a_i$ [8]. In other words the only “accented” symbol occurring in U_i is a_i , i.e., $U_i \in (A \cup \{\hat{a}_i\})^*$.

Associated to U_i is a $\{0, 1\}$ -word $\nu(U_i) = \nu_K \nu_{K-1} \cdots \nu_k$ where $\nu_K = 10$, $\nu_j = \varepsilon$ (the empty word) if $\sigma(x_j) \neq a_i$, $\nu_j = 10$ if $\sigma(x_j) = a_i$ and $m_j = \rho(x_j)$, $\nu_j = 010$ if $\sigma(x_j) = a_i$ and $0 < m_j < \rho(x_j)$ and $\nu_j = 00$ if $\sigma(x_j) = a_i$ and $m_j = 0$.

By comparing the matrix formulation given in Corollary 2.11 in [8] used to compute $R_\omega(U_i)$ with the matrix formulation given in Proposition 4.1 in [1], we leave it to the reader to verify the following:

PROPOSITION 3.1. — $R_\omega(U_i) = R_2(\nu(U_i))$.

In other words computing the multiplicities of representations in a generalized Ostrowski numeration system may be reduced to a computation of the multiplicities of representations in the Fibonacci base.

Example 3.2. — We consider the example originally started in Berstel’s paper [1] and later revisited by Justin and Pirillo as Example 2.3 in [8] of the Ostrowski numeration system associated to the infinite word $\omega = a, a, b, b, a, a, a, b, b, a, a, b, b, a, a, a, b, \dots$. It is readily verified that

$$\mathcal{N}(\omega) = \{1, 3, 7, 24, 55, 134, 323, \dots\},$$

$\sigma(1) = \sigma(7) = \sigma(55) = \sigma(323) = a$, $\sigma(3) = \sigma(24) = \sigma(134) = b$, and $\rho(1) = 2$, $\rho(3) = 2$, $\rho(7) = 3$, $\rho(24) = 2$, $\rho(55) = 2$, $\rho(134) = 2$, $\rho(323) = 3$.

Applying the greedy algorithm we obtain the following representation of the number 660

$$660 = 2(323) + 0(134) + 0(55) + 0(24) + 2(7) + 0(3) + 0(1).$$

So $Z_\omega(660) = \hat{a}\hat{a}bbaabb\hat{a}\hat{a}bbaa$. which is a semigood multiblock based at a . We deduce that

$$\nu(Z_\omega(660)) = 10 \cdot \varepsilon \cdot 00 \cdot \varepsilon \cdot 010 \cdot \varepsilon \cdot 00$$

or simply $\nu(Z_\omega(660)) = 100001000$.

Following the algorithm of Corollary 2.11 of [8] due to Justin and Pirillo, we obtain $q_1 = 2$, $q_2 = 4$, $p_1 = 2$, $p_2 = 2$, $c_1 = c_2 = 1$, so that

$$R_\omega(660) = (1, 0) \begin{pmatrix} 0 & 2 \\ 0 & 3 \end{pmatrix} \begin{pmatrix} 0 & 2 \\ 0 & 3 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 6$$

In contrast, applying the algorithm in Proposition 4.1 of [1] due to Berstel to the Zeckendorf word $\nu(Z_\omega(660)) = 100001000$, we obtain $d_1 = 4$, $d_2 = 3$ so that

$$R_2(\nu(Z_\omega(660))) = (1, 1) \begin{pmatrix} 1 & 1 \\ 2 & 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = 6$$

as required⁽²⁾

Acknowledgements

The second author was partially supported by a grant from the National Security Agency.

BIBLIOGRAPHY

- [1] J. BERSTEL, "An exercise on Fibonacci representations, A tribute to Aldo de Luca", *RAIRO, Theor. Inform. Appl.* **35** (2002), p. 491-498.
- [2] V. BERTHÉ, "Autour du système de numération d'Ostrowski", *Bull. Belg. Math. Soc. Simon Stevin* **8** (2001), p. 209-239, Journées Montoises d'Informatique Théorique (Marne-la-Vallée, 2000).
- [3] L. CARLITZ, "Fibonacci representations", *Fibonacci Quarterly* **6(4)** (1968), p. 193-220.
- [4] N. FINE & H. WILF, "Uniqueness theorem for periodic functions", *Proc. Amer. Math. Soc.* **16** (1965), p. 109-114.
- [5] O. JENKINSON & L. ZAMBONI, "Characterizations of balanced words via orderings", *Theoret. Comput. Sci.* **310** (2004), p. 247-271.

⁽²⁾In [1], Berstel computes $R_\omega(660)$ in a different way by using the matrix formulation of Proposition 5.1 in [1] which applies to an Ostrowski numeration system.

- [6] J. JUSTIN, “Algebraic combinatorics and Computer Science”, chap. Episturmian words and morphisms (results and conjectures), p. 533-539, Springer Italia, Milan, 2001.
- [7] J. JUSTIN & G. PIRILLO, “Episturmian words and Episturmian morphisms”, *Theoret. Comput. Sci.* **302** (2003), p. 1-34.
- [8] ———, “Episturmian words: shifts, morphisms and numeration systems”, *Internat. J. Found. Comput. Sci.* **15** (2004), p. 329-348.
- [9] J. JUSTIN & L. VUILLON, “Return words in Sturmian and Episturmian words”, *Theor. Inform. Appl.* **34** (2000), p. 343-356.
- [10] P. KOCÁBOVÁ, Z. MASÁCOVÁ & E. PELANTOVÁ, “Ambiguity in the m -bonacci numeration system”, preprint, 2004.
- [11] A. OSTROWSKI, “Bemerkungen zur Theorie der Diophantischen Approximation I”, *Abh. Math. Sem. Hamburg* **1** (1922), p. 77-98.
- [12] R. THJEDMAN & L. ZAMBONI, “Fine and Wilf words for any periods”, *Indag. Math. (N.S.)* **14** (2003), p. 135-147.
- [13] E. ZECKENDORFF, “Représentation des nombres naturels par une somme de nombres de Fibonacci ou de nombres de Lucas”, *Bull. Soc. Royale Sci. Liège* **42** (1972), p. 179-182.

Marcia EDSON & Luca Q. ZAMBONI
University of North Texas
Department of Mathematics
PO Box 311430
Denton, TX 76203-1430 (USA)
mre0006@unt.edu
luca@unt.edu